

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 11/00, 11/02, 11/08, 5/00	A1	(11) International Publication Number: WO 98/51785 (43) International Publication Date: 19 November 1998 (19.11.98)
(21) International Application Number: PCT/US97/08254 (22) International Filing Date: 14 May 1997 (14.05.97) (71) Applicants: THE GENERAL HOSPITAL CORPORATION [US/US]; 55 Fruit Street, Boston, MA 02114 (US). MASSACHUSETTS INSTITUTE OF TECHNOLOGY [US/US]; 77 Massachusetts Avenue, Cambridge, MA 02142-1324 (US). (72) Inventors: BHATIA, Sangeeta; 92 Cottage Street, Sharon, MA 02067 (US). YARMUSH, Martin; Apartment 403, 950 Massachusetts Avenue, Cambridge, MA 02139 (US). TONER, Mehmet; 100 Pilgrim Road, Wellesley, MA 02181 (US). (74) Agent: MEIKLEJOHN, Anita, L.; Fish & Richardson, P.C., 225 Franklin Street, Boston, MA 02110-2804 (US).		(81) Designated State: JP. Published <i>With international search report.</i>
(54) Title: CO-CULTIVATION OF CELLS IN A MICROPATTERNED CONFIGURATION		
(57) Abstract <p>Methods are provided for producing co-cultures of cells in which at least two types of cells are configured in a micropattern on a substrate such that modulating of the extent of heterotypic cell-cell contacts can be used to modulate metabolic and/or synthetic functions of cells. Co-culture can be carried out by coating a protein such as collagen on a substrate so that the protein defines a micropattern on the substrate, contacting the coated substrate with cells of a first type such as hepatocytes in a medium to bind the cells to the protein to produce a micropatterned cell-coated substrate, contacting the micropatterned substrate with cells of a second type such as fibroblasts in a second medium to bind the cells to the substrate. The first or second medium is a non-attachment medium such as a serum-free medium and the other medium is an attachment medium such as a medium containing serum. The attachment medium may contain an attachment factor such as fibronectin or an antibody. The micropattern may be formed by islands of cells of either the first or second type surrounded by cells of the other type.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakistan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

- 1 -

CO-CULTIVATION OF CELLS IN A MICROPATTERNED CONFIGURATIONStatement as to Federally Sponsored Research

This invention was produced, at least in part,
5 with funds from the United States Government under
National Institutes of Health Grant DK5270. Therefore,
the United States Government has certain rights in the
invention.

Background of the Invention

10 The invention relates to methods for co-
cultivating cells in micropatterned formations (e.g., for
the production of bioartificial organs).

Co-cultures of hepatocytes with another cell type
have been recognized to prolong cell survival rates,
15 maintain phenotype and induce albumin secretion in
hepatocytes. Heretofore, such co-cultures have been
investigated by other researchers, but not in a
configuration that allows for the manipulation or control
of the interaction of the two cell types in the culture.
20 In preparation of such co-cultures, cells of one type are
seeded to a substrate and are allowed to attach for a
time until cells of a second type are seeded atop the
cells of the first type. In such co-cultures, parameters
such as cell number are controllable, but they lack the
25 definition and control of the spatial orientation.
Heterotypic cell-cell contacts are left to random and the
ability to upregulate or optimize certain cell functions
is not possible in such cultures (Clement, B., et al.
"Long-Term Co-Culture of Adult Human Hepatocytes with Rat
30 Liver Epithelial Cells: Modulation of Albumin Secretion
and Accumulation of Extracellular Material" Hepatology
4(3): 373-380 (1984); Schrode, W., et al. "Induction of
Glutamine Synthetase in Periportal Hepatocytes by
Cocultivation with a Liver Epithelial Cell Line"
35 European Journal of Cell Biology 53: 35-41 (1990);
Michalopoulos, G., et al. "Primary Culture of

- 2 -

Hepatocytes on Human Fibroblasts" In Vitro 15(10):
796-806 (1979); Guguen-Guillouzo, C., et al.
"Maintenance and Reversibility of Active Albumin
Secretion by Adult Rat Hepatocytes Co-Cultured with
5 Another Liver Epithelial Cell Type" Experimental Cell
Research 143: 47-54 (1983); Begue, J. et al. "Prolonged
Maintenance of Active Cytochrome P-450 in Adult Rat
Hepatocytes Co-Cultured with Another Liver Cell Type"
Hepatology 4(5): 839-842 (1984); Agius, L. "Metabolic
10 Interactions of Parenchymal Hepatocytes and Dividing
Epithelial Cells in Co-culture" Biochemical Journal 252:
23-28 (1988); and Reid, L. et al. "Culturing Hepatocytes
and Other Differentiated Cells" Hepatology 4(3): 548-559
(1984)).

15

Summary of the Invention

The invention provides methods for producing co-
cultures of cells in which at least two types of cells
are configured in a micropattern on a substrate. By
20 using micropatterning techniques to modulate the extent
of heterotypic cell-cell contacts, it is now possible to
modulate (e.g., upregulate or downregulate) metabolic
and/or synthetic functions of cells.

Accordingly, the invention provides a method for
25 producing a micropatterned co-culture containing at least
two cell types; the method entails:

- i) providing a protein-coated substrate, wherein
a protein coating the substrate defines a micropattern on
the substrate;
- 30 ii) contacting the protein-coated substrate with
cells of a first cell type suspended in a first cell
medium under conditions such that cells of the first cell
type bind the protein of the protein-coated substrate,
thereby producing a micropatterned cell-coated substrate;
- 35 and

- 3 -

iii) contacting the micropatterned cell-coated substrate with cells of a second cell type suspended in a second cell medium under conditions such that cells of the second cell type bind the substrate, thereby
5 producing the micropatterned co-culture, wherein one of the cell media is a non-attachment medium and one of the cell media is an attachment medium.

Typically, in practicing the invention, the cells of the first and second cell types are mammalian cells,
10 although the cells may be from two different species (e.g., humans, rats, mice, etc). Examples of suitable combinations of cells for producing the co-culture include, without limitation:

- a) hepatocytes (e.g., primary hepatocytes) and
15 fibroblasts (e.g., NIH 3T3-J2 cells);
- b) hepatocytes and at least one other cell type, particularly liver cells, such as Kupffer cells, Ito cells, endothelial cells, and biliary ductal cells;
- c) endothelial cells and smooth muscle cells;
- 20 d) tumorigenic parenchymal cells and mesenchymal cells;
- e) hematopoietic cells and bone marrow cells (e.g., adipocytes, fibroblasts); and
- f) skin cells (e.g., keratinocytes) and
25 fibroblasts. Other combinations of cells also are within the invention.

The substrate on which the cells are grown can be any biologically compatible material to which cells can adhere, such as glass, polymers (such as fluoropolymers,
30 fluorinated ethylene propylene, polyvinylidene, polydimethylsiloxane, polystyrene, polycarbonate, and polyvinyl chloride), and silicon substrates (such as fused silica, polysilicon, or single silicon crystals). To produce a micropattern of the co-cultured cell types,
35 protein is first adhered to the substrate in order to

- 4 -

define (i.e., produce) a micropattern. The micropattern produced by the protein serves as a "template" for formation of the cellular micropattern. Typically, a single protein will be adhered to the substrate, although
5 two or more proteins may be used to define the micropattern (for example, one micropatterned protein may be used to attract one cell type, while a second micropatterned protein is used to attract a second cell type). In practicing the invention, a variety of
10 techniques can be used to foster selective cell adhesion of two or more cell types to the substrate. Included, without limitation, are methods such as localized protein adsorption, organosilane surface modification, alkane thiol self-assembled monolayer surface modification, wet
15 and dry etching techniques for creating three-dimensional substrates, radiofrequency modification, and ion-implantation (Lom, 1993; Brittlund, 1992; Singhvi, 1994b; Singhvi, 1994a; Ranier, 1994; Bellamkonda, 1993; and Valentini, 1993).

20 Proteins that are suitable for producing a micropattern are those proteins to which one of the cell types of the co-culture specifically binds under the cell culture conditions used to cultivate the co-culture (i.e., conventional cell culture conditions). For
25 example, hepatocytes are known to bind to collagen. Therefore, collagen is well-suited to facilitate binding of hepatocytes in a micropattern. Other suitable proteins include fibronectin, laminin, and entactin, and combinations thereof.

30 Typically, in practicing the invention, the cells of the first cell type (e.g., hepatocytes) initially are suspended in an "non-attachment" cell culture medium (e.g., serum-free medium and media that lack "attachment factors"), while the cells of the second cell type are
35 suspended in an "attachment" medium [e.g., a cell culture

- 5 -

medium that contains serum (typically 1-10% (e.g., 5-10%)), or one or more "attachment factors" (typically at least 1 ng/ml (e.g., 5-100 ng/ml)) such as fibronectins and other extracellular matrix, selectins, RGD peptides, ICAMs, E-cadherins, and antibodies that specifically bind a cell surface protein (for example, an integrin, ICAM, selectin, or E-cadherin)]. In another method of practicing of the invention, the cells of the second type have intrinsic attachment capabilities, thus eliminating a need for the addition of serum or exogenous attachment factors. Some cell types will attach to electrically charged cell culture substrates and will adhere to the substrate via cell surface proteins and by secretion of extracellular matrix molecules. Fibroblasts are an example of one cell type that will attach to cell culture substrates under these conditions. Thus, the invention also includes a method for producing a micropatterned co-culture containing at least two cell types where the method entails:

- i) providing a protein coated substrate wherein a protein coated substrate defines a micropattern on the substrate;
- ii) contacting the protein-coated substrate with cells of a first cell type suspended in a first cell medium under conditions such that the cells of the first cell type bind the protein of the protein-coated substrate, thereby producing a micropatterned cell-coated substrate; and
- iii) contacting the micropatterned cell-coated substrate with cells of a second cell type suspended in a second cell medium under conditions such that the cells of the second cell type bind to the substrate, thereby producing the micropatterned co-culture, wherein the first cell type (e.g., dermal fibroblasts of skin) is in non-attachment medium and second cell type has natural

- 6 -

attachment capabilities to attach it to the substrate. A charged substrate is particularly useful in practicing this variation of the invention.

In a variation of these methods for producing
5 micropatterned co-cultures, the first cell type can be genetically engineered (using conventional techniques) to produce a desired gene product, and the second cell type produces a protein, and the first cell type enables the second cell type to reproduce and grow.

10 By using micropatterning techniques, such as those described herein, the first and second cell types define a micropattern (i.e., are configured into a pattern having a resolution on a micron scale). In the micropattern of the co-culture, cells of either the first
15 or second cell type are surrounded by cells of either the second or first cell type, respectively. For example, the cells of the co-culture can be configured such that "islands" of hepatocytes (cells of a first cell type) are surrounded by fibroblasts (cells of a second cell type).
20 Such islands need not be perfectly circular in shape. For, example, the islands can be produced as stripes or rectangles. Regardless of the shape of the island, it is preferred that at least 30% of the cells of the island are within 100 μm of an interface between the island of
25 cells (e.g., hepatocytes) and the surrounding cells (e.g., fibroblasts). More preferably, at least 50%, 80% or 90% of the cells of the island are within 100 μm of the interface. Where the island is essentially circular, the island typically will have a diameter of 25-1,000 μm
30 (preferably, 30-500 μm (or 100-500 μm)).

In a variation of the above methods, the invention provides a method for upregulating a metabolic or synthetic function of a cell of a first cell type; the method entails:

- 7 -

i) providing a protein-coated substrate, wherein a protein coating the substrate defines a micropattern on the substrate;

ii) contacting the protein-coated substrate with
5 cells of a first cell type suspended in a first cell medium under conditions such that cells of the first cell type bind the protein of the protein-coated substrate, thereby producing a micropatterned cell-coated substrate; and

10 iii) contacting the micropatterned cell-coated substrate with cells of a second cell type suspended in a second cell medium under conditions such that cells of the second cell type bind the substrate, thereby producing the micropatterned co-culture, wherein:

15 a) one of the cell media is a non-attachment medium and one of the cell media is an attachment medium; and

b) the cells of the first and second cell types define a micropattern wherein cells of the second cell
20 type surround cells of the first cell type, and at least 30% of the cells of the first cell type are within 100 μm of an interface between the cells of the first cell type and the cells of the second cell type,

thereby producing a micropatterned co-culture,
25 wherein a metabolic or synthetic function of a cell of the first cell type is upregulated relative to cells of the first cell type in an unpatterned co-culture that comprises cells of the first and second cell types.

This method derives from our observations that by
30 using micropatterning techniques to modulate the level of heterotypic cell-cell contact in a co-culture, one can upregulate a synthetic or metabolic function of a cell in the co-culture. For example, DNA and/or protein synthesis can be upregulated with this micropatterning
35 method. In a micropatterned co-culture where islands of

- 8 -

hepatocytes are surrounded by fibroblasts, the upregulation of cell function can be detected as an increase in intracellular or secreted albumin of a hepatocyte. Alternatively, or in addition, upregulation
5 of cell function can be detected as an increase in urea synthesis in a hepatocyte.

As in the above-described methods for co-cultivating cells in a micropatterned configuration, examples of suitable combinations of cells for the co-
10 culture include, without limitation,

- a) hepatocytes (e.g., primary hepatocytes) and fibroblasts (e.g., NIH 3T3-J2 cells);
- b) hepatocytes and at least one other cell type, particularly liver cells, such as Kupffer cells, Ito
15 cells, endothelial cells, and biliary ductal cells;
- c) endothelial cells and smooth muscle cells;
- d) tumorigenic parenchymal cells and mesenchymal cells;
- e) hematopoietic cells and bone marrow cells
20 (e.g., adipocytes, fibroblasts); and
- f) skin cells (e.g., keratinocytes) and fibroblasts.

Referring to the above list, the invention typically will be practiced such that an island of the first-named cell
25 type in each of these combinations is surrounded by cells of the second-named cell type, and the function of the first-named cell type is upregulated. In producing the micropatterned co-culture, it is not necessary to adhere to the substrate the cell whose function is to be
30 upregulated prior to adhering the other cell. However, when producing a co-culture of hepatocytes and fibroblasts, the hepatocytes typically will be adhered to the protein-coated substrate prior to contacting the substrate with the fibroblasts. Other parameters of this
35 aspect of the invention (e.g., island size, attachment

- 9 -

factors, substrate, etc.) are essentially as described above.

Typically, the metabolic and/or synthetic function of cells of the first cell type is modulated at least 1.5-fold in micropatterned co-cultures, relative to a metabolic or synthetic function of cells of the first cell type in an unpatterned co-culture. As shown by the experiments described below, an increase of at least 5-10-fold also is achievable. To detect the modulation of a metabolic or synthetic function, conventional molecular and biochemical assays can be used, such as those described below.

In practicing this method, we have found that, not only is cell function upregulated to a higher absolute level (e.g., of albumin production) in the micropatterned co-cultures (relative to unpatterned co-cultures), but also the kinetics of this upregulation are increased. In other words, the rate at which a metabolic or synthetic function is upregulated to a particular level in the micropatterned co-culture is increased relative to the rate at which a metabolic or synthetic function is upregulated in an unpatterned co-culture. Thus, the invention also provides a method for modulating the kinetics at which metabolic or synthetic functions of a cell are upregulated in a co-culture. From a bioengineering perspective, this increase in the kinetic of cell function upregulation is advantageous as it decreases the cultivation time needed for cells to reach a particular level of metabolic or synthetic function. In practice, we have found that an unpatterned co-culture may take 1-2 weeks to reach a particular level of cell function, whereas a micropatterned co-culture could be upregulated to that level in a single day.

Also included within the invention are the micropatterned co-cultures produced according to the

- 10 -

methods described herein. Such micropatterned co-cultures of cells can be used as bioartificial organs for *in vivo*, *ex vivo*, or *in vitro* purposes. For example, a micropatterned co-culture of hepatocytes combined with
5 fibroblasts can be used as an implantable (*in vivo*) or extracorporeal (*ex vivo*) artificial liver for replacement of liver function (e.g., in response to diseases, infections, or trauma), or in *in vitro* assays of liver
10 function (for example, for toxicology or basic research purposes). Similarly, such micropatterned co-cultures can be used as a source of protein (e.g., albumin or clotting factors from hepatocytes). In this regard, the invention provides the advantage over cell-free methods of producing proteins because intracellular post-
15 translational modifications that occur in the co-cultures of the invention will provide a properly modified (e.g., glycosylated) protein.

As used herein, the term "micropattern" refers to a pattern formed on a substrate (e.g., by a protein,
20 cell, or combination of cells of two or more types), which has a spatial resolution (e.g., 1-5 μm) that permits spatially controlling cell placement at the single-cell level. Thus, using micropatterning methods, one can precisely manipulate cell-cell interactions. In
25 contrast, in an "unpatterned" co-culture of cells, the cells are randomly distributed.

As used herein, an "island" of cells is a single cell, or typically a group of cells, of one cell type that is surrounded by cells of another cell type (e.g., a
30 group of hepatocytes surrounded by fibroblasts). Thus, an interface is formed where cells at the periphery of the island meet the surrounding cells. An island need not be circular in shape; for example, rectangular islands, and islands of other, amorphous shapes can be
35 used in the invention. Regardless of the shape of the

- 11 -

island, however, at least 30% (preferably at least 50%, 80%, or 90%) of the cells in the island should be within 100 μm from an interface between the cell types. Thus, where the island is essentially circular in shape, islands that are less than 1,000 μm in diameter are suitable. Typically, the island will be 30-500 μm in diameter.

Brief Description of the Drawings

Fig. 1 is a schematic representation of a process to generate micropatterned co-cultures.

Figs. 2A-2D are a series of photographs obtained by immunofluorescent staining of micropatterned co-cultures. Fig. 2A shows micropatterned co-cultures as detected by immunofluorescence of cytokeratin. Fig. 2B shows randomly-distributed co-cultures as detected by immunofluorescence of cytokeratin. Fig. 2C shows micropatterned co-cultures as detected by immunofluorescence of F-actin. Fig. 2D shows randomly-distributed co-cultures as detected by immunofluorescence of F-actin.

Fig. 3 is a schematic representation of a method for determining X, the heterotypic interaction parameter.

Fig. 4 is a schematic representation of a method to obtain separation of cell populations.

Description of the Preferred Embodiments

The working examples provided herein are to illustrate, not limit, the invention. Various parameters of the scientific methods employed in these examples are described in detail below and provide guidance for practicing the invention in general.

In these particular working examples, hepatocytes are co-cultured with fibroblasts; as is described herein, similar methods can be used to co-culture other combinations of cells. These experiments demonstrate, for the first time, the micropatterning of two cell types

- 12 -

in a co-culture. In other words, the two cell types can be used to define a pattern having a resolution on a micron scale. These experiments also show that, by increasing the extent of heterotypic cell-cell contacts in the co-culture, the metabolic and synthetic functions of cells of the micropatterned co-culture are upregulated, relative to cells in an unpatterned configuration.

PART I

10 MATERIALS AND METHODS

Microfabrication techniques were used to modify glass substrates with biomolecules. These modified substrates were utilized to pattern a single cell type or micropattern co-cultures in various configurations. Fig. 15 1 schematically depicts the overall process for producing micropatterned co-cultures.

Microfabrication of Substrates

The experimental substrates were produced utilizing standard microfabrication techniques. Chrome masks of the desired dimensions were generated on a pattern generator (Gyrex), which transferred the pattern to a chromium coated quartz plate using a contact printer and a developer. Round, 2" diameter X 0.02" thickness borosilicate wafers (Erie Scientific) were cleaned in a piranha solution (3:1 H₂SO₄: 30% H₂O₂) for 10 minutes, rinsed, and blown dry with a N₂ gun. Wafers were then dehydrated by baking for 60 minutes at 200°C. Discs were subsequently coated with positive photoresist (OCG 820-27 centistokes) on a Headway spin-coater with vacuum chuck as follows: dispense photoresist at 500 RPM for 2 seconds, spread photoresist at 750 RPM for 6 seconds, spin at 4000 RPM for 30 seconds, resulting in a 1 μm coating (Step A, Fig. 1). Wafers were then pre-baked for 5 minutes at 90°C to remove residual solvent and anneal any stress in the film. Wafers were exposed in a

- 13 -

Bottom Side Mask Aligner (Karl Suss) to ultraviolet light through the desired chromium mask to create a latent image in the resist layer. Exposure occurred under vacuum-enhanced contact for 3 seconds. Exposed photoresist was then developed to produce the final three-dimensional relief image for 70 seconds in developer (OCG 934 1:1), rinsed three times under running deionized water and cascade rinsed for 2 minutes (Step B, Fig. 1). Subsequently, discs were hard-baked for 30 minutes at 120°C to remove residual developing solvents and promote adhesion of the film. Finally, substrates were exposed to oxygen plasma at 250 W for 4 minutes to remove unwanted resist in areas to be subsequently modified. Wafers were stored at room temperature for up to 2 months. Substrates were subsequently re-exposed to oxygen plasma 24 hours prior to further processing to ensure availability of borosilicate for surface modification on a Plasma Day Etcher at a base vacuum of 50 mTorr and O₂ pressure of 100 mTorr at a power of 100 W for 2-4 minutes.

Surface Modification of Substrates

Substrates were modified using experimental methods similar to those developed by Lom et al. and Britland et al. (Step C, Fig. 1) (Stenger et al., 1992; Lom et al, 1993). Briefly, substrates were rinsed twice in distilled, deionized (DD) water and allowed to air dry. Silane immobilization onto exposed glass was performed by immersing samples for 30 seconds in freshly prepared, 2% v/v solution of 3-[(2-aminoethyl)amino] propyltrimethoxysilane (AS, Hüls America) in water followed by 2 rinses in 200 mL DD water. Wafers were then dried with nitrogen gas and baked at 120 °C for 10 minutes. Next, discs were immersed in 20 mL of 2.5% v/v solution of glutaraldehyde in PBS (pH 7.4) for 1 hour at 25 °C. Substrates were then rinsed twice in fresh PBS,

- 14 -

and immersed in a 4 mL solution of a 1:1 solution of 1 mg/mL collagen I (Dunn et al, 1991): DD water for 15 minutes at 25°C. Discs were subsequently immersed in acetone and placed in a bath sonicator (Bransonic) for 15 minutes to remove residual photoresist ultrasonically (Step D, Fig. 1). Wafers were then rinsed twice in DD water, and soaked overnight in 70% ethanol for sterilization (Step E, Fig. 1).

Surface Characterization of Substrates

10 Autofluorescence. Wafers were observed using a Nikon Diaphot microscope equipped with a Hg lamp and power supply (Nikon). The autofluorescence of photoresist (excitation: 550 nm, emission: 575 nm) was used to visualize micropatterned substrates prior to surface modification. Absence of autofluorescence after 15 sonication was taken to verify indicate removal.

Profilometry. Profilometry was performed to characterize surface topology at the Center for Material Science Engineering (CMSE) on a Dektak 3 Profilometer 20 (Veeco Instruments) with a 12.5 μm radius probe at a scan rate of 100 $\mu\text{m/s}$.

Atomic Force Microscopy (AFM). AFM was performed in order to characterize the spatial distribution of immobilized groups at the CMSE, MIT on a Nanoscope 3 25 (Digital Instruments) equipped with a standard 117 μm silicon cantilever operating in tapping mode with a scan size of 100 μm .

Indirect Immunofluorescence of Collagen I.

Collagen-derivatized substrates were incubated at 37°C 30 with undiluted Rabbit Anti-Rat Collagen I Antisera (Biosciences) by inverting substrates onto parafilm that contained a droplet (50 μL) of antisera for 1 hour. Substrates were then washed thoroughly in PBS and placed on a rotating shaker at 25°C for 30 minutes. This 35 washing procedure was repeated twice. Next, discs were

- 15 -

inverted onto parafilm with 50 μ L (1:20) of Dichlorotriazinylamino Fluorescein (DTAF)-conjugated Donkey Anti-Rabbit IgG (Jackson) in blocking solution. Blocking solution consisted of 3% w/w bovine serum albumin, 1% donkey serum, 0.04 % sodium azide, pH 7.4. Finally, substrates were washed in PBS overnight, and observed by fluorescence microscopy (excitation: 470 nm, emission: 510 nm).

Cell Culture

10 Hepatocyte Isolation and Culture. Hepatocytes were isolated from 2- to 3-month-old adult female Lewis rats (Charles River) weighing 180-200 g, by a modified procedure of Seglen (1976). Detailed procedures for isolation and purification of hepatocytes were previously described by Dunn et al (1989). Routinely, 200-300 million cells were isolated with viability between 85% and 95%, as judged by Trypan blue exclusion. Non-parenchymal cells, as judged by their size (< 10 μ m in diameter) and morphology (nonpolygonal or stellate), were less than one percent. Culture medium was Dulbecco's modified Eagle's Medium (DMEM, Gibco) supplemented with 10% fetal bovine serum (FBS, JR Scientific), 0.5 U/mL insulin, 7 ng/mL glucagon, 20 ng/mL epidermal growth factor, 7.5 (g/mL hydrocortisone, 200 U/mL penicillin , 200 (g/mL streptomycin and 50 (g/mL gentamycin ('hepatocyte media with serum'). Serum-free culture medium was identical except for the inclusion of 40 (g/mL of L-Proline (Sigma) and exclusion of FBS (Lee et al, 1993) ('serum-free hepatocyte media').

30 NIH 3T3-J2 Culture. NIH 3T3-J2 cells were provided by Howard Green (Harvard Medical School). Cells grown to pre-confluence were trypsinized in 0.01% trypsin (ICN Biomedicals)/0.01% EDTA (Boehringer Mannheim) solution in PBS for 5 minutes and then resuspended in 25

- 16 -

mL media. Approximately 10% of the cells were inoculated into a fresh tissue culture flask. Cells were passaged at pre-confluency no more than 12 times. Cells were cultured in 75 cm³ flasks (Corning) at 10% CO₂, balance moist air. Culture medium consisted of DMEM (Gibco) with high glucose, supplemented with 10% bovine calf serum (BCS, JRH Biosciences) and 200 U/mL penicillin and 200 µg/mL streptomycin.

Cell Culture on Modified Surfaces. Wafers were rinsed in sterile water, and incubated in 0.05 % w/w bovine serum albumin in water at 37°C for 1 h to pre-coat substrates with a non-adhesive protein. Substrates were then washed twice with serum-free media. Next, hepatocytes were seeded at high density (4×10^6 /mL) in serum-free media for 1.5 hours at 37 °C, 10% CO₂, balance air (Step E, Fig. 1). Surfaces were then rinsed twice by pipetting and then aspirating 4 mL of serum-free media, re-seeded with hepatocytes for 1.5 hours, rinsed with 4 mL of serum-free media, and incubated overnight (Step F, Fig. 1). The following day, 3T3 cells were trypsinized as described above, counted with a hemocytometer and plated at 1×10^6 /mL in 2 mL of serum-containing, high glucose DMEM, and allowed to attach overnight (Step G, Fig. 1). 'Randomly-distributed' (i.e., unpatterned) co-cultures consisted of hepatocyte seeding in the desired number (usually 250,000) on a uniformly collagen-derivatized surface followed by 3T3 seeding after 24 hours.

Immunofluorescent Staining

Cultures were washed 2 times with 2 mL PBS, fixed and permeabilized with 10 mL of acetone at -20 °C for 2 minutes, and washed twice in 10 mL PBS. Cultures on wafers were incubated at 37 °C with undiluted Rabbit Anti-Rat Pan Cytokeratin Antisera (Accurate Chemical), by inverting substrates onto parafilm containing a 50 µL

- 17 -

droplet of antisera for 1 hour. Substrates were then washed, incubated with secondary antibody, and washed (as described above for indirect immunofluorescence of collagen). Secondary antibody also included

5 rhodamine-phalloidin (1:100, Molecular Probes) for fluorescent labeling of F-actin. Specimens were observed and recorded using a Nikon Diaphot microscope (Nikon) equipped with a Hg lamp and power supply (Nikon), light shuttering system (Uniblitz D122), CCD camera

10 (Optitronics CCD V1470), and MetaMorph Image Analysis System (Universal Imaging) for digital image acquisition.

Image Analysis

To quantitatively describe the extent of heterotypic interactions we measured the fraction of cell

15 perimeter in contact with adjacent cells of a different cell type (X). For example, $X=1$ for a single cell island whereas $X=0$ for a cell amidst hepatocyte neighbors. Images were acquired as described above and analyzed with MetaMorph Image Analysis System. Cells were sampled from

20 each field and manually outlined to obtain individual cell perimeters, P. Subsequently, the regions of heterotypic cell-cell contact were similarly delineated, F. Each cell was assigned its characteristic $X = F/P$ and these values of X were averaged over 20-50 cells for each

25 condition. For population distributions, individual values of X were assigned to an appropriate 'bin', and histograms plots were generated.

RESULTS

As is discussed below, we performed surface

30 characterization studies on substrates in the absence of cells to first verify the ability to obtain spatially-defined surface chemistries. Subsequently, the ability to micropattern single cell cultures and

- 18 -

co-cultures including two different cell types was shown, as is described below.

Surface Characterization

Topological and spatial uniformity of photoresist patterns were assessed using profilometry and autofluorescent properties of photoresist. The photoresist coating was found to be approximately $1.35 \mu\text{m}$ in thickness using the specified spin-coating parameters. Furthermore, the thickness of photoresist varied $<5\%$ within each scan. Autofluorescence of photoresist was utilized to examine integrity and distribution of photoresist prior to and during processing. Autofluorescent regions corresponding to $\sim 1 \mu\text{m}$ variations in thickness were detected. Absence of any contaminant fluorescence in the dark lanes indicates complete, uniform removal of exposed photoresist during development.

In order to verify regional aminosilane (AS) modification of borosilicate, substrates were exposed to AS followed by removal of photoresist. Aminosilane modification has been previously reported to modify the three-phase contact angle of water with the surface (Lom et al, 1993); therefore, the perimeter of a single water droplet should display microscopic undulations on patterns of varying hydrophilicity. These undulations were observed; $20 \mu\text{m}$ AS modified lanes exhibit differential wetting properties relative to the adjacent $50 \mu\text{m}$ unmodified lanes. Therefore, selective AS modification of exposed glass was verified in the pattern of the original $20 \mu\text{m}/50 \mu\text{m}$ striped photoresist pattern, indicating the ability for photoresist to serve as a 'chemical mask' to AS modification of underlying glass.

- 19 -

Collagen immobilization via glutaraldehyde derivatization of patterned AS surfaces was also characterized.

Fluorescence micrographs were obtained, showing the results of indirect immunofluorescent staining of areas of presumed collagen immobilization. Fluorescent regions, corresponding to regions of collagen localization, were patterned uniformly with spatial resolution on the micron level. Furthermore, fluorescent patterns corresponded to initial photoresist patterns without evidence of undercutting. More importantly, despite processing in acetone and 70% ethanol, collagen retained sufficient immunoreactivity for antibody binding.

Collagen-derivatized surfaces were also analyzed with AFM to determine differences in topology between unmodified and modified borosilicate. Modified regions with a width of 20 μm were found to have an average height of 4 nm above the unmodified, 50 μm lanes. These data can be utilized to approximate the number of collagen monolayers atop AS.

Micropatterning of Co-Cultures

The aforementioned experiments demonstrate the ability to reproducibly utilize photoresist patterns to generate immobilized collagen patterns; the following experiments illustrate the applicability of these techniques to cellular micropatterning. Seeding of the first cell type, hepatocytes, resulted in localization to collagen-derivatized regions and normal polygonal morphology. The cellular configurations were dictated by the positioning of collagen on glass which was in turn controlled by the choice of chromium mask in the microfabrication procedure. Furthermore, hepatocytes conformed to the edges of the collagen pattern on the modified glass. Typical hepatocyte diameter in suspension is 20 μm , whereas, upon attachment and

- 20 -

unconstrained spreading, cell diameters increase to 30-40 microns. Therefore, after attachment to 20 μm lines, cells were observed to elongate in the axial direction upon spreading. Similar cytoskeletal changes were
5 observed in cells on corners of larger patterns or on the perimeter of circular patterns.

The versatility of this technique was seen in phase-contrast micrographs. Initial hepatocyte patterns of 20 μm and 200 μm were modified by the addition of
10 fibroblasts in serum-containing media. Fibroblasts were observed to localize solely to unmodified (glass) regions of patterned substrates resulting in micropatterned co-cultures of 20 μm /50 μm and 200 μm /500 μm . The grating pattern utilized was chosen for its illustrative
15 potential; however, in principle, co-cultures can be achieved in any desired configuration. Thus, our approach is clearly adaptable to both micropatterning of single cell cultures and co-cultures of two different cell types.

20 Spreading of the primary cell type typically resulted in negligible residual sites of collagen-derivatization. Therefore, attachment of the secondary cell type in this example should be limited either to unmodified glass or the surface of the primary
25 cell type. We determined that 3T3 fibroblasts do not undergo significant attachment to hepatocyte surfaces by performing plating experiments of fibroblasts on monolayers of hepatocytes which showed no attachment even after a 4 hour incubation (data not shown). In addition,
30 fibroblast attachment and spreading on glass was characterized by seeding cells in serum-containing media on glass coverslips where they were observed to attach and spread with high efficiency within 4 hours (data not shown).

- 21 -

Indirect immunofluorescence was utilized to selectively stain cell populations and aid in visual discrimination between different cell types. We compared presence of cytokeratin (Figs. 2A and 2B), an intermediate filament expressed in hepatocytes but absent in mesenchymal cells, to F-actin (Figs. 2C and 2D), a cytoskeletal protein present in all mammalian cells. We also produced a comparison of a patterned co-culture of 200 μm /500 μm , compared to a 'randomly distributed' co-culture with identical attached cell numbers of both cell populations. Of note, was the level of homotypic hepatocyte interaction in Fig. 2A, a 200 μm , stripe of cells versus Fig. 2B, a random distribution of cells. Hepatocytes in the 200 μm stripe had primarily homotypic contacts, whereas those in Fig. 2B had predominantly heterotypic contacts. Furthermore, the distribution of heterotypic interaction over a patterned cell population is observed to be greatly reduced over random co-cultures, where hepatocytes are shown to be present in single cell islands, doublets, and triplets (Fig. 2B).

To quantitatively describe the extent of heterotypic contact, we used image analysis and perimeter tracing to define the fractional cell perimeter engaged in heterotypic cell contact as X , as described above. Fig. 3 schematically depicts sample perimeter tracings (black lines) with high-lighted interfaces of heterotypic contacts corresponding to hepatocytes in a digitally-acquired phase micrograph. This particular pattern (200 μm /500 μm) has very little heterotypic contact, as was visually observed; therefore, the average X over the population is small due to the majority of cells with $X=0$. We demonstrated the ability to vary the mean value of X over a cell population from 0.7 in the randomly distributed culture to 0.08 using

- 22 -

micropatterning. Moreover, different patterns (20/50) produce distinct mean values of X ($X = 0.55$). Variations of X from the mean were also examined for randomly distributed cultures as compared to defined
5 patterns (20/50). As observed microscopically, hepatocytes in randomly distributed cultures experience heterogeneous microenvironments - single hepatocytes, doublets, and multicellular aggregates can be observed within a given culture. Quantitative analysis of
10 population distributions corroborate the variability in X in randomly distributed cultures as compared to micropatterns (20/50 and 50/50), which exhibited a relatively small variance around the mean value of X . Thus, variations in cellular microenvironment, both in
15 amount and variability, were achieved without varying the numbers of cells in each sub-population.

DISCUSSION

Many conventional co-culture systems have been limited by the inability to vary cell local seeding
20 density independently of the cell number as well as inherent variations in the distribution of cell contacts over a population of cells. We have developed a versatile technique for the micropatterning of two different cell types derived from conventional strategies
25 for surface modification with aminosilanes linked to biomolecules and by manipulating the serum content of cell culture media, as described above. This co-culture technique allows the manipulation of the initial cellular microenvironment without variation of adhered cell
30 number. Specifically, we were able to control both the degree and type of initial cell-cell contact. Differences in homotypic and heterotypic interaction were demonstrated, allowing variations in exposure to cell-surface receptors, locally secreted extracellular
35 matrix, and local concentrations of soluble factors.

- 23 -

The patterning methodology utilized in this particular working example departs from certain conventional methods in several ways. For example, AS was applied after photoresist patterning but before photoresist lift off. This differs from Lom et al (1993), who apply AS before photoresist application. In these systems, incubation with biomolecules, such as horseradish peroxidase, results in adsorption to both regions which becomes specific after denaturation of non-specifically bound protein in 8M urea. Many proteins, such as collagen, undergo irreversible adsorption (Deyme et al, 1986) and will therefore not desorb from unmodified regions. We preserved the integrity of the photoresist throughout the surface modification process and removed the photoresist after the deposition of collagen. This was achieved by deposition of AS in water, which does not normally attack photoresist. AS is known to oligomerize in aqueous solution (Arkles et al, 1991), but is stable at least for a period of hours. In this way, we utilized photoresist to mask the borosilicate from non-specific protein adsorption and did not need to rely on protein denaturation and desorption, nor on AS deposition prior to photoresist patterning.

Atomic force microscopy was utilized to approximate the depth of the immobilized collagen layer. Modified regions were ~4 nm above the unmodified regions. AS molecules have been estimated to have a height of 1.2 nm end-to-end (Lom et al, 1993). In the helical configuration, collagen I fibrils have dimensions of 300 nm in length and 1.2 nm in diameter (Darnell et al, 1990). These data suggest that there were 1-2 layers of collagen fibrils, configured lengthwise, corresponding to an upper limit of 0.1 $\mu\text{g}/\text{cm}^2$ per monolayer of 'side-on' packed fibrils (Deyme et al, 1986). Therefore,

- 24 -

achievable collagen surface concentrations are within an order of magnitude those observed in adsorbed collagen systems ($0.37 \mu\text{g}/\text{cm}^2$) (Deyme et al, 1986). Another consideration is the bioactivity of biomolecules after exposure to acetone and ethanol. We have demonstrated preservation of bioactivity of collagen I via cell attachment and spreading as well as by antibody binding for indirect immunofluorescence. Others have shown functional integrity of laminin, fibronectin, collagen IV, and bovine serum albumin (Lom et al, 1993). Proteins sensitive to acetone may benefit from adaptation of the photoresist lift-off procedure.

In this particular example, selective cell attachment to immobilized collagen was enhanced by adsorption of bovine serum albumin to the surface to deter non-specific adhesion. This has been shown to be effective for hepatocytes, neuroblastoma cells (Lom et al, 1993; Miyamoto et al, 1993; Bhatia et al, 1994), and many antibody-antigen interactions. In contrast, albumin adsorption to (mono)aminosilane on fluorinated ethylene propylene films has been shown to have the opposite effect of mediating attachment of aortic endothelial cells (Ranieri et al, 1993), thus, this aspect of this particular procedure may be modified for optimal results when using other cell types.

Using primary rat hepatocytes and 3T3 fibroblasts, we demonstrated the ability to vary initial heterotypic (X) interactions over a wide range, while preserving the ratio of cell populations in culture. Thus, co-culture interactions now can be manipulated in an entirely new dimension. In addition, micro-patterned co-cultures were observed to have less variation in the level of heterotypic contacts (X) than did random co-cultures. Therefore, measurement of macroscopic biochemical quantities in micro-patterned co-cultures will provide

- 25 -

better representations of specific cell-cell interactions than those seen in unpatterned co-cultures.

In general, this methodology for micropatterning co-cultures can also be applied to other techniques of patterning biomolecules, such as self-assembled monolayers. In addition, three-phase co-cultures can be established by patterning of two different, cell-specific biomolecules.

SUMMARY

The invention provides a simple, versatile technique for controlling homotypic versus heterotypic interactions of at least two cell types in culture. One can vary X without changing the number of cells in each sub-population and therefore vary the ratio of cell types in a given culture.

PART II

The following experiments demonstrate that the methods of the invention can be used to upregulate metabolic and synthetic functions (e.g., liver-specific functions) of cells in the micropatterned co-culture.

MATERIALS AND METHODS

Substrates for micropattern formation were prepared essentially as described above.

Hepatocyte isolation and culture

Hepatocytes were isolated from 2- to 3-month-old adult female Lewis rats (Charles River Laboratories, Wilmington, MA) weighing 180-200 g, by a modified procedure of Seglen (1976). Detailed procedures for isolation and purification of hepatocytes were previously described by Dunn et al (FASEB, 1989). Routinely, 200-300 million cells were isolated with viability between 85% and 95%, as judged by trypan blue exclusion. Nonparenchymal cells, as judged by their size (<10 μ m in diameter) and morphology (nonpolygonal or stellate), were less than 1%. Culture medium was Dulbecco's modified

- 26 -

eagle's medium (DMEM, Gibco) supplemented with 10% fetal bovine serum (FBS, Sigma, St.Louis, MO), 0.5 U/mL insulin, 7 ng/mL glucagon, 20 ng/mL epidermal growth factor, 7.5 mg/mL hydrocortisone, 200 U/mL penicillin, and 200 mg/mL streptomycin ('hepatocyte media with serum'). Serum-free culture medium was identical except for the exclusion of FBS.

NIH 3T3-J2 Culture

NIH 3T3-J2 cells were provided by Howard Green (Harvard Medical School). Cells grown to preconfluence were passaged by trypsinization in 0.01% trypsin (ICN Biomedicals, Costa Mesa, CA)/0.01% EDTA (Boehringer Mannheim, Indianapolis, IN) solution in PBS for 5 minutes, diluted, and then inoculated into a fresh tissue culture flask. Cells were passaged at pre-confluency no more than 10 times. Cells were cultured in 175 cm² flasks (Fisher, Springfield, NJ) at 10% CO₂, balance moist air. Culture medium consisted of DMEM (Gibco, Grand Island, NY) with high glucose, supplemented with 10% bovine calf serum (BCS, JRH Biosciences, Lenexa, KS) and 200 U/mL penicillin and 200 mg/mL streptomycin ('fibroblast media'). In some cases, growth arrested cells were obtained for DNA analysis by incubation with 10 mg/mL mitomycin C (Boehringer Mannheim) in media for 2 hours (reconstituted just prior to use) followed by three washes with media. Mitomycin C-treated fibroblasts were shown to have constant levels of DNA for 10 days of culture, verifying the lack of fibroblast growth under these conditions.

Cell culture on modified surfaces

Wafers were sterilized by soaking for 2 hours in 70% ethanol in water at room temperature. Subsequently, wafers were rinsed in sterile water and incubated in 0.05% w/w bovine serum albumin (BSA) in water at 37°C for 1 hour to precoat substrates with a nonadhesive protein.

- 27 -

Substrates were then placed in sterile P-60 tissue culture dishes (Corning, Corning, NY), and rinsed in sterile water followed by a final rinse with serum-free media. Next, hepatocytes were seeded at high density
5 (1-2 x 10⁶/mL) in 2 mL serum-free media for 1.5 hours at 37°C, 10% CO₂, balance air followed by two rinses with serum-free media. This process was repeated twice to ensure confluence of hepatocytes, especially on larger dimension patterns. The following day, 3T3 cells were
10 trypsinized as described above, counted with a hemocytometer, plated at 3.75 x 10⁵/mL in 2 mL of serum-containing, high-glucose DMEM, and allowed to attach overnight. Subsequently, 2 mL hepatocyte culture media (described above) was sampled and replenished
15 daily.

Experimental Design

Spatial configurations of micropatterned co-cultures were manipulated by varying mask dimensions. Transparent circular areas (or 'holes') on chrome masks
20 correspond to derivatized, and ultimately hepatocyte-adhesive, areas of glass substrates. In order to achieve identical hepatocyte numbers across varying micropatterned configurations, the total surface area of all 'holes' was kept constant across all masks despite
25 changes in hole diameter and center-to-center spacing. All arrays were hexagonally packed with the exception of the largest dimension hole which consisted of a single unit of 17800 μm diameter. Thus, pattern dimensions varied as follows (hole diameter (in microns),
30 center-to-center spacing (in microns)): 36, 90; 100, 250; 490, 1229; 6800, 16900; and a single unit of 17800 μm diameter, where the resulting total hepatocyte-adhesive area on 2" diameter glass substrates would be identical in all cases.

- 28 -

Analytical Assays

Media samples were collected daily and stored at 4°C for subsequent analysis for urea and albumin content. Urea synthesis was assayed using a commercially available
5 kit (Sigma Chemical Co., kit No. 535-A). Reaction with diacetyl monoxime under acid and heat yields a color change detected at 540 nm. Albumin content was measured
10 by enzyme-linked immunosorbent assays (ELISA) as described previously (Dunn et al, 1991). Rat albumin and anti-rat albumin antibodies were purchased from Cappel
Laboratories (Cochranville, PA).

DNA analysis was adapted from a method of MacDonald and Pitt (1991). Cells were sacrificed at the indicated time of culture by washing with PBS, removal
15 and immersion of wafer into PBS to eliminate dead cells underneath the substrate, and subsequent incubation with 0.05% (w/v) type 4 collagenase (Sigma) in Kreb's Ringer Buffer at 37° C for 30 minutes to release the cell layer from the underlying substrate. Next, cells were removed
20 with a rubber policeman and the cell/collagenase mixture was removed. The substrate was washed with PBS which was then combined with the above solution. The resulting solution was combined with an equivalent volume of hepatocyte media for neutralization of collagenase,
25 followed by centrifugation at 1000 RPM for 5 minutes. The supernatant was aspirated, and cells were resuspended in 2 mL PBS. Subsequently, the samples were frozen at -80°C for up to 1 month.

For analysis, the frozen samples were rapidly
30 thawed in a 37° C water bath to promote membrane rupture. Freeze-thaw protocols have been established as an effective way to rupture the cell membrane in order to gain access to cellular contents (Rago et al, 1990). To ensure complete cell lysis, samples were then sonicated
35 using a probe sonicator (Branson) for 10 seconds at 4°C.

- 29 -

Samples were vortexed and 20 ml samples were placed into a 96-well plate (NUNC, Denmark). Similarly, 20 ml of DNA standard (double stranded Calf Thymus DNA, Sigma) in PBS from 100 to 0 mg/mL were vortexed and placed on each
5 plate. This volume was combined with 100 ml salt/dye buffer (2 M NaCl, 10 mM Tris, 1 mM EDTA, 1.6 mM Hoechst 33258 (Molecular Probes, Eugene, OR)). Samples and standards were allowed to incubate with salt/dye buffer at room temperature in the dark for 30 minutes before
10 reading on a Spectrofluorometer (Millipore, Bedford, MA) Excitation 360 nm, 1/2 bandwidth 40 nm, Emission, 460 nm, 1/2 bandwidth 40 nm.

Analysis of DNA Content

The total DNA content in cultures with
15 growth-arrested fibroblasts was assayed as follows. Mitomycin C was utilized to growth arrest fibroblasts (as described above) and 1.5×10^6 fibroblasts were counted with a hemocytometer and added to micropatterned hepatocyte cultures. Replicate cultures were either
20 sacrificed 6 hours after fibroblast seeding or after 9 days of co-culture and assayed for total DNA as described above.

Immunohistochemistry

Cultures were washed twice with PBS, fixed with 4
25 % paraformaldehyde in PBS for 30 minutes, and permeabilized for 10 minutes with 0.1% Triton in PBS. Endogenous avidin-binding activity of hepatic tissue was blocked by 20 minute incubations with 0.1% avidin and 0.01% biotin in 50 mM Tris-HCl respectively (Biotin
30 Blocking System X590, DAKO, Carpinteria, CA). Endogenous peroxidase activity was blocked by 30 minute incubation with a hydroxgen peroxide and sodium azide solution (Peroxidase Blocking Reagent, DAKO). Rabbit anti-rat albumin antibodies (Cappell) were utilized with
35 horse-radish peroxidase visualization by use of a

- 30 -

biotinylated secondary antibody to rabbit IgG,
streptavidin-labelled horse radish peroxidase, and
hydrogen peroxide in the presence of
3-amino-9-ethylcarbazole as a substrate (Rabbit Primary
5 Universal Peroxidase Kit #K684, DAKO).

Functional Bile Duct Staining

Cultures were washed three times with media and
incubated for 5 hours with 2 μ M Carboxyfluorescein
diacetate (Molecular Probes) in an adapted method of
10 LeCluyse et al. (1994). Subsequently, cultures were
washed again three times and examined microscopically.
Digital images were obtained on a Nikon Diaphot
microscope equipped with Hg lamp and excited at 470 nm
excitation and 510 nm emission.

15 Image Acquisition and Analysis

Specimens were observed and recorded using a Nikon
Diaphot microscope equipped with a CCD camera (Optronics
CCD V1470), and MetaMorph Image Analysis System
(Universal Imaging, Westchester, PA) for digital image
20 acquisition. Image analysis on immunostained images was
performed utilizing the thresholding function in
MetaMorph and visual correlation with brightfield images.

Statistics and Data Analysis

Experiments were repeated two to three times with
25 duplicate or triplicate culture plates for each
condition. Two duplicate wells were measured for
biochemical analysis. One representative experiment is
presented where the same trends were seen in multiple
trials but absolute rates of production varied with each
30 animal isolation. Each data point represents the mean of
three dishes. Error bars represent standard error of the
mean. Statistical significance was determined using
one-way ANOVA (analysis of variance) on Statistica
(StatSoft) with Tukey HSD (Honest Significant Difference)
35 Post-Hoc analysis with $p < 0.05$.

- 31 -

RESULTS

Micropatterned co-cultures were generated with variations in heterotypic interface yet identical surface area (i.e., cell numbers) dedicated to both hepatocyte and fibroblast adhesion. Five different configurations ranging from maximal heterotypic contact (smallest islands) to minimal heterotypic contact (single island) were characterized for expression of liver-specific function by use of: two biochemical markers (albumin and urea synthesis), immunohistochemistry (intracellular albumin staining), transport across apical surface (bile duct excretion), and DNA content. These results show an increase in heterotypic interactions is correlated with an increase in liver-specific functions.

15 Characterization of Initial Cell Distribution

All 5 micropatterns were designed to have similar levels of hepatocyte-adhesive surface area (2.5 cm^2), which is expected to correspond to identical number of attached hepatocytes. Variations in spatial configurations were utilized to generate differences in total perimeter of hepatocyte islands from 5.6 cm to 2800 cm, which, upon addition of fibroblasts, should correspond to variations in the total heterotypic interface. Micropatterns ranged from many single hepatocyte islands of $36 \mu\text{m}$ diameter to a single island of 17.8 mm diameter ($100 \mu\text{m}$, $490 \mu\text{m}$, and $6800 \mu\text{m}$ islands also were detected). Micropatterned hepatocytes were found to adhere predominantly to collagen-modified areas in all 5 conditions with close agreement between theoretical and observed values for total initial hepatocyte island perimeter (data not shown).

To verify similar numbers of attached hepatocytes across various spatial configurations, we measured DNA content of micropatterned hepatocyte cultures 24 hours after hepatocyte seeding (i.e. prior to fibroblast

- 32 -

seeding). All cultures had statistically similar levels of DNA ($8 \pm 1.8 \mu\text{g}$) with the exception of increased DNA content ($18 \pm 3.3 \mu\text{g}$) on the smallest island ($36 \mu\text{m}$ diameter) micropatterns.

5 The smallest islands were designed to produce single cell islands. The dimensions of these islands ($36 \mu\text{m}$ diameter) was chosen to correspond with the experimentally determined projected surface area of a single, spread hepatocyte on immobilized collagen I of
10 $1000 \mu\text{m}^2$ (data not shown); however, isolated rat hepatocytes have a diameter of approximately $20 \mu\text{m}$, allowing the potential for individual islands to retain more than one hepatocyte upon seeding with a concentrated cell suspension. In addition, hepatocytes have been
15 shown to have an increased mitotic index at low seeding densities (Nakamura et al, 1983), which may have contributed to increased hepatocyte DNA in this condition. To distinguish between increased cell number as compared to increased ploidy, image analysis of one
20 thousand $36 \mu\text{m}$ micropatterned islands was completed at 6 hours after initiation of cell seeding. This analyses demonstrated more than one cell per island in 57% of cases, with an average of 1.9 ± 1.2 cells per island. Therefore, we concluded that increased DNA was due to
25 increased hepatocyte number on the smallest pattern.

 Addition of 3T3-J2 fibroblasts to micropatterned hepatocytes resulted in micropatterned co-cultures with marked alterations in initial heterotypic interface despite similar numbers of fibroblasts and hepatocytes
30 across conditions. Phase contrast micrographs of 4 of the 5 configurations ($36 \mu\text{m}$, $100 \mu\text{m}$, $480 \mu\text{m}$, and $6800 \mu\text{m}$ islands) demonstrated the significant variation in hepatocyte microenvironment which was achieved by altering micropattern dimensions.

- 33 -

Biochemical Analysis of Liver-Specific Function

To demonstrate the effect of modulation of the local hepatocyte environment on liver-specific function, albumin secretion and urea synthesis were measured as markers of differentiated function. These two markers were measured as a function of micropattern dimensions in the presence and absence of fibroblasts. In cultures of fibroblasts alone, albumin secretion and urea synthesis by fibroblasts was found to be undetectable; therefore changes in these markers in co-cultures were attributed to differences in hepatocyte metabolism.

Albumin secretion for five different spatial configurations was determined for pure hepatocyte cultures. A rapid decline in liver-specific functions was detected for all five conditions (36 μm , 100 μm , 490 μm , 6800 μm , and 17,800 μm islands), from initial values of $8.8 \pm 0.9 \mu\text{g/day}$ to undetectable levels.

We also measured albumin secretion for the same five micropatterns with the addition of fibroblasts. Albumin synthesis increased over time in culture in all configurations from less than 10 $\mu\text{g/day}$ to greater than 34 $\mu\text{g/day}$, indicating up-regulation of this liver-specific function due to co-culture with fibroblasts. These micropatterned co-cultures had decreasing amounts of initial heterotypic contact with maximal levels occurring at the smallest hepatocyte island dimension (36 μm) and minimal levels occurring at the single large hepatocyte island (17.8 mm). Smaller islands with high levels of heterotypic contact demonstrated greater albumin secretion than larger islands (less heterotypic contact) after day 5 of culture. Two fundamental patterns of up-regulation were observed: (1) dramatic up-regulation to similar levels of albumin secretion in the three smallest island configurations (19 to 26-fold of initial levels) and (2)

- 34 -

relatively modest up-regulation (~7-fold) in the two larger island configurations. Therefore, a three-fold increase in albumin production was achieved in certain pattern configurations by modulation of the initial cellular microenvironment.

Analysis of urea synthesis in micropatterned co-cultures revealed similar qualitative results. Urea synthesis was either constant over culture or increased from less than ~100 $\mu\text{g/day}$ to 160 $\mu\text{g/day}$ indicating up-regulation of another liver-specific function due to co-cultivation with fibroblasts. In addition, two patterns of up-regulation were observed using this marker of differentiated function: (1) up-regulation of urea synthesis to similar levels in the three smallest island configurations (up to 2-fold increase), and (2) relatively little up-regulation in the two larger island configurations. Therefore, a two-fold increase in urea synthesis production was achieved in certain pattern configurations by modulation of the initial cellular microenvironment. Asterisks indicate $p < 0.05$ in Tukey post-hoc analysis of variance.

Hepatocyte Function In Situ: Immunostaining of Intracellular Albumin

In order to further examine the observed variations in liver-specific function exhibited by various micropatterned co-cultures, we studied the hepatocyte phenotype in situ by immunostaining of intracellular albumin. Specifically, we first focused on the distribution of albumin staining as it related to the heterotypic interface in one representative pattern, 490 μm hepatocyte islands (at days 2 and 6). In addition, in order to distinguish between homotypic effects on differentiation and the effects arising from varying the heterotypic interface, we performed immunostaining on micropatterned pure hepatocyte cultures

- 35 -

at days 2 and 6. Our results show that hepatocytes cultured alone stained uniformly for intracellular albumin at 48 hours after isolation. The level of protein declined subsequently on the order of days. In
5 comparison, micropatterned co-cultures displayed a more complex behavior. They, too, displayed initial uniform staining for intracellular albumin. Over 6 days, however, hepatocytes close to the heterotypic interface stained for high levels of intracellular albumin, whereas
10 protein levels in hepatocytes far from the heterotypic interface ($> 3-4$ cells) continued to decline as in the pure hepatocyte cultures. To ensure that this 'ring' of intense staining was due to variations in intracellular albumin content of hepatocytes, as opposed to the
15 detachment of hepatocytes or fibroblasts from the lightly-stained areas, phase contrast microscopy of these cultures was performed. The presence of fibroblasts in the periphery of the hepatocyte island and cellular structures in the center of the hepatocyte island was
20 clearly depicted. This peripheral 'ring' of intense staining observed across a $490\ \mu\text{m}$ micropatterned co-culture was reproducible.

In order to correlate the pattern of immunostaining with the variations that we observed by
25 biochemical analysis of secreted products in media, we also examined the distribution of high levels of intracellular albumin in comparatively small ($100\ \mu\text{m}$) and large ($6800\ \mu\text{m}$) micropatterned co-cultures. These micrographs demonstrate uniform intense staining in
30 smaller islands (initial island size $100\ \mu\text{m}$), a well-demarcated ring of $\sim 120\ \mu\text{m}$ in intermediate size islands (initial size $490\ \mu\text{m}$), and a well-demarcated ring of $\sim 380\ \mu\text{m}$ in larger islands (initial size $6800\ \mu\text{m}$), indicating a negative correlation between differentiated

- 36 -

hepatocyte phenotype and distance from the heterotypic interface.

Hepatocyte Function in Situ: Bile Duct Excretion

Another in situ marker of liver-specific function
5 is the formation of functional bile caniliculi between
hepatocytes. Carboxyfluorescein diacetate (CFDA) is
taken up by hepatocytes, cleaved by intracellular
esterases, and in the presence of normal biliary
transport, excreted across the apical membrane of the
10 hepatocyte. The presence of normal biliary transport of
the dye as well as functional integrity of the
tight-junctional domains bounding the caniliculus, causes
illumination of visibly fluorescent bile duct structures
between hepatocytes. We probed two patterns, one from a
15 highly functioning co-culture (490 μm circles) and one
from a poorly functioning group (17800 μm circle), as
determined by albumin and urea production, in order to
examine this marker of liver-specific function. Phase
contrast micrographs of both cultures were produced. We
20 found that 490 μm patterns developed functional bile
caniliculi, especially in the island periphery, whereas
fluorescent bile duct staining was not observed on 17800
 μm islands.

DISCUSSION

25 In this set of experiments, we demonstrated that
liver-specific tissue function can be modulated by
controlling initial heterotypic cell-cell interactions,
despite use of identical cellular components.
Furthermore, it was determined that these differences in
30 bulk tissue properties as a function of cellular
microenvironment were generated by induction of spatial
heterogeneity in the hepatocyte phenotype. Hepatocytes
in the vicinity of the heterotypic interface had a
relative increase in levels of liver-specific function;

- 37 -

therefore, spatial configurations with maximal initial interface exhibited superior function.

Cellular Microenvironment Modulated Liver-specific Functions

5 Evidence that liver-specific function could be controlled by variations in initial cell-cell interactions is seen in the functional differences between predominantly heterotypic co-cultures (smallest islands of 36 μm diameter) and predominantly homotypic
10 co-cultures (largest island of 17800 μm diameter) as assessed by markers of metabolism (urea synthesis), synthetic function (albumin secretion and cytoplasmic content), and apical transport (biliary excretion). These cellular microenvironments significantly altered
15 liver-specific functions as follows: increasing hepatocyte island size correlated with a relative decline in urea synthesis, albumin secretion, intracellular albumin staining, and effective biliary excretion. Smaller hepatocyte islands of 36, 100, and 490 μm initial
20 diameter yielded three-fold steady-state increases in albumin secretion and two-fold steady-state increases in urea synthesis over 6800 and 17800 μm islands. Similarly, a smaller pattern (490 μm initial diameter) exhibited functional biliary excretion as assessed by
25 accumulation of a fluorescent compound within bile canilicular structures between hepatocytes whereas larger islands (17,800 μm initial diameter) showed reduced functional biliary excretion with no evidence of focal fluorescence. The presence of fluorescent biliary
30 structures between hepatocytes was correlated by LeCluyse et al. (1994) to biliary structures observed on electron microscopic analysis. The absence of fluorescent biliary structures was attributed to either (1) low rate of excretion across apical domain (2) absence or loss of

- 38 -

function of tight junctions at borders of apical membrane or (3) decreased uptake of dye by hepatocytes. The lack of fluorescent biliary structures in 17800 μ m pattern indicates some such functional deficit. Therefore,

5 hepatocytes in smaller island co-cultures have improved biliary transport as well as relative improvements in other liver-specific functions due to alterations in the initial cellular microenvironment (as compared with larger island co-cultures).

10 In concluding that bulk tissue function (secreted albumin and urea) was modulated by initial cellular microenvironment, hepatocyte numbers were measured to ensure that changes in these liver-specific markers were due to changes in level of hepatocellular function (as
15 opposed to differences in cell division). In order to assess the relative contribution of hepatocyte division, as compared to up-regulation of functions, fibroblasts were growth-arrested, and total DNA in co-cultures was measured. Thus, changes in total DNA could be attributed
20 solely to hepatocytes. Total DNA of co-cultures was measured at 6 hours of co-culture and compared to DNA content at 9 days of co-culture. This analysis demonstrated that no significant increase in total DNA occurred in co-cultures over 9 days, indicating increases
25 in hepatic functions were due to up-regulation of synthesis rather than a marked increase in hepatocyte population (data not shown). These data correlate well with reports of minimal hepatocyte division under various co-culture configurations (Guguen-Guillouzo, 1986;
30 Kuri-Harcuch and Mendoza-Figuera, 1989; Donato et al, 1990). Furthermore, this result correlated well with visual observation of larger micropatterns (490 micron island diameter and greater), where hepatocyte island size was observed to be relatively constant over the
35 course of culture, indicating a lack of significant cell

- 39 -

division. Taken together, these data indicate that variations in hepatic functions between culture configurations were due predominantly to relative levels of hepatic upregulation, as opposed to hepatocyte
5 division.

The conclusion that bulk tissue function was modulated by variation of the cell-cell interactions at the heterotypic interface prompted confirmation of similar initial hepatocyte numbers to confirm that
10 changes in secreted products were due to up-regulation of liver-specific functions, rather than differences in numbers of initial hepatocytes. Comparison of initial total hepatocyte DNA in all five micropatterns showed this to be a valid approximation ($8 \pm 1.8 \mu\text{g}$ DNA) with,
15 perhaps, the exception of the smallest ($36 \mu\text{m}$) islands which were found to have two-fold elevated levels of DNA. This may be due to the potential for more than one unspread hepatocyte ($20 \mu\text{m}$ diameter) to adhere to $36 \mu\text{m}$ islands. In any event, in these studies, the trend to
20 increased long-term liver specific function resulting from maximal initial heterotypic interface remained a consistent finding.

The experiments described above were conducted with the same surface area dedicated to fibroblasts in
25 all conditions. This allowed examination of the local cellular environment as an isolated variable, without differences in cell numbers and resultant variations in concentrations of potential signaling factors (such as humoral factors in media). In addition, these
30 experiments allowed simultaneous control over both oxygen delivery to hepatocytes, as well as amount of media. In contrast, variation of culture plate area necessitates either a change in media volume to preserve the depth of media above the cell population (and the diffusion of
35 oxygen) or a change in media depth to preserve media

- 40 -

volume. Therefore, these methods for controlling cellular environment have definitively demonstrated the importance of local cellular microenvironment as an isolated modulator of liver-specific function (i.e.,
5 metabolic and synthetic functions).

Cellular Microenvironment Induced Spatial Heterogeneity
in Hepatocyte Phenotype

In addition to demonstrating that liver-specific tissue function can be modulated by controlling initial
10 heterotypic cell-cell interactions, the experiments described herein demonstrate that spatial heterogeneity in the induction of the hepatocyte phenotype was the primary cause of these variations in function. In situ immunostaining of intracellular albumin on micropatterned
15 hepatocyte/fibroblast co-cultures displayed increased staining in the vicinity of the heterotypic interface, indicating up-regulation of this marker of differentiated function. Specifically, smaller (100 μm islands) stained throughout hepatocyte regions whereas larger islands (490
20 μm and greater) exhibited intense staining in a well-demarcated ring in the periphery. This pattern of staining was highly reproducible both spatially and across various conditions. The differentiated hepatocyte phenotype appeared to dominate within 100-400 μm of the
25 heterotypic interface; therefore, it is reasonable to conclude that patterns with greater interfacial regions displayed superior tissue function.

To confirm that variations in intracellular albumin represented variations in hepatocyte phenotype
30 due to heterotypic interactions, the effect of homotypic hepatocyte interactions on the spatial distribution of intracellular albumin in a representative micropattern (490 μm island) was assessed. These experiments revealed uniform staining in pure hepatocyte cultures with

- 41 -

decreased staining over a period of one week, consistent with the observed decline in secreted albumin and previous studies showing residual albumin mRNA hepatocyte immediately after isolation with decline of mRNA over 1
5 week (Dunn et al, 1992); therefore, patterns of immunostaining in co-cultures were indeed due to heterotypic interactions with fibroblasts, rather than homotypic interactions.

Image Analysis: In order to correlate
10 intracellular albumin staining with albumin secretion data, image analysis was performed on immunostained co-cultures. Specifically, we estimated the fraction of hepatocytes contributing to albumin secreted into the media. Image analysis of intracellular albumin staining
15 revealed ~100% of hepatocytes stained intensely in 100 μ m patterns, ~65% in 490 μ m patterns, and ~20% in 6800 μ m patterns. By assuming negligible contribution of weakly staining hepatocytes to albumin production, we calculated that hepatocytes adjacent to the heterotypic interface in
20 larger patterns may have produced 35-50% more albumin per cell than those in 100 μ m micropatterns. These data suggest there may be a further increase in albumin production in hepatocytes adjacent to relatively undifferentiated homotypic neighbors.

25 Micropatterning co-cultures, as described above, allowed the creation of larger hepatocyte colonies than those that come about by random aggregation and cell migration; therefore, these assays was able to demonstrate a finite penetration length of a
30 differentiation signal to the interior of a large hepatocyte colony. This result contradicts the notion that hepatocytes are able to communicate effectively throughout a hepatocyte colony.

- 42 -

Related Observations on Control of Cell-Cell Interactions

While the ability to micropattern co-cultures provides the ability to modulate tissue function via the initial cellular microenvironment, the inherent dynamics of cell adhesion and motility may further modify these engineered tissues. The degree of morphogenesis depended upon hepatocyte island size. In these experiments, hepatocyte islands of 490 μm with center-to-center spacing of 1230 μm produced a relatively stable configuration whereas hepatocytes in islands of 100 μm and smaller underwent some reorganization into cord-like structures. Reorganization of tissue may be prevented by cytoskeletal toxins such as cytochalasin D. Despite the tendency for some spatial configurations to reorganize, the perturbations which were achieved in initial cellular microenvironment were found to have significant long-term impact on tissue function.

SUMMARY

These experiments show that microfabrication can be used as a vehicle to control heterotypic cell-cell interactions without significant variations in cell numbers. Unprecedented modulation of heterotypic interface as an independent variable was achieved. This modulation of the heterotypic interface over three orders of magnitude dramatically altered levels of detectable liver-specific function in the resulting composite tissues as measured by markers of metabolic, synthetic, and excretory function. Variations in function were due to modulation of the hepatocyte phenotype: specifically, epithelial differentiation varied inversely with distance from the heterotypic interface, causing cultures with a relative increase in cell interaction to exhibit superior function. The ability to control heterotypic cell-cell interactions and probe the resulting tissue for evidence

- 43 -

of cell communication has applications both in basic science (e.g., *in vitro* assays of tissue function) and development of functional tissue constructs for medical applications. From a fundamental perspective, these co-
5 culture techniques can be exploited in assays for determining the mechanisms by which cells communicate. In the area of tissue engineering, the ability to co-cultivate two or more cell types in a micropattern and modulate cell function provides an unprecedented level of
10 control over the *in vitro* reconstruction of skin, bone marrow, muscle, and many other tissues.

PART III

In the following experiments, we used microfabrication techniques as well as conventional
15 culture methodologies to further examine the mechanism of induction of hepatocyte differentiation at the heterotypic interface. These experiments indicate that the biological signal for the observed induction of hepatic functions is "cell-associated" (broadly defined
20 to include membrane-bound receptors, locally secreted extracellular matrix, and local matrix or cell-bound growth factors), rather than "freely secreted" (broadly defined to include humoral factors such as soluble cytokines and growth factors).

25 MATERIALS AND METHODS

Examination of the modes of cell communication in hepatocyte/3T3 co-culture was conducted using *in situ* immunostaining to assess the contribution of homotypic hepatocyte interactions, and various methods of probing
30 the class of signal(s) responsible for induction of the hepatocyte phenotype in hepatocytes proximal to the heterotypic interface. These techniques included pre-treated media to probe for soluble factors (conditioned media), separation of cell populations to
35 probe for labile soluble factors and to eliminate

- 44 -

contribution of fibroblast adhering to the hepatocyte surface (spacer), and cultures conducted with continual disturbance of overlying media to probe for transport limitations (agitation).

5 General Techniques

Methodology for micropatterned substrate preparation, hepatocyte isolation and culture, NIH 3T3-J2 fibroblast culture, immunohistochemistry, analytical assays, and image acquisition are presented in detail
10 above.

Immunostaining of Micropatterned Cultures

To assess the contribution of hepatocyte homotypic interaction on spatial patterns of albumin immunostaining, various sizes of micropatterned
15 hepatocytes were probed both in the presence and absence of additional fibroblasts. Micropatterned cultures of hepatocytes alone and hepatocyte/fibroblast co-cultures were generated as described above in the following hepatocyte island dimensions: 36, 100, 490, 6800, and
20 17800 μm . Hepatocytes were either cultured alone or co-cultured with 750,000 NIH 3T3-J2 fibroblasts. Culture media (2 mL) was replaced daily. Cultures were fixed and stained at 48 hours and 144 hours.

Conditioned Media

25 Conditioned media experiments were performed in unpatterned configurations. Glass substrates were modified by aminosilane, glutaraldehyde, and collagen I as described in above, resulting in collagen I immobilization over the entire wafer. Hepatocytes were
30 seeded in 'hepatocyte media with serum' as described previously, at a density of 250,000 per P-60. Four different culture configurations were investigated. First, in order to control for baseline degradation of biochemical compounds in media at 37°C, hepatocytes were

- 45 -

fed daily with 2 mL of media which had been previously incubated for 24 hours in tissue culture plastic. Second, in order to examine the effects of fibroblast secreted products, hepatocytes were fed daily with 2 mL of media which had previously incubated for 24 hours with (750,000 initially seeded) NIH 3T3-J2 cells on an unmodified glass wafer. Third, in order to probe the effects of fibroblast secreted products which require hepatocyte interaction for their up-regulation, hepatocytes were fed daily with 2 mL of media which had been previously incubated for 24 hours with a co-culture of (750,000 initially seeded) NIH 3T3-J2 cells and 250,000 hepatocytes on an, unpatterned, collagen-modified wafer. Last, in order to generate a 'positive control' to compare the above conditions to co-culture induced up-regulation of liver-specific functions, hepatocytes were co-cultured with NIH 3T3-J2 fibroblasts by the addition of 750,000 NIH 3T3-J2 cells on day 2 of culture. Media was collected daily and stored at 4°C.

Physical separation of cell types

Hepatocytes and fibroblasts were separated by the following general protocol: placement of a polymer annulus on glass substrate, surface modification of glass within the annulus by adsorption of collagen I, attachment of hepatocytes to central, collagen-immobilized region, 'capping' of hepatocyte population during fibroblast seeding to prevent access of fibroblasts to top surface of hepatocytes, and removal of cap and annulus. Differential spacing was achieved by variation in annulus width resulting identical inner diameter (and therefore size of hepatocyte island) and larger outer diameter (resulting in larger separation between cell populations). Fig. 4 depicts a schematic overview of method.

- 46 -

Annuli were fabricated with polydimethylsiloxane (PDMS) (Sylgard 184, Dow Corning, Lansing, MI). Stock sheets of 500 μ m thickness were prepared by casting polymer solution (mixed as described by the manufacturer) in polystyrene tissue culture plastic for 2 hours at 65°C. Annuli were fabricated with inner diameter of 0.6 cm and various outer diameters using disposable skin punch biopsy cutting tools. To limit potential cytotoxicity, PDMS annuli were then coupled to collagen I with aminoethylaminopropyltrimethoxysilane and glutaraldehyde as previously described.

'Caps' were fabricated from sheets of polyethylene terephthalate (PET) by use of a standard paperpunch to generate 0.6 cm disks from 7 mil thickness mylar film (Kodak). Discs were soaked in 70% ethanol in water for 2 hours followed by rinsing in media.

Annuli were affixed to clean, 2" diameter, borosilicate wafers, and subsequently 'heat-fixed' to prevent detachment via three consecutive exposures to a heat gun at a distance of 10 cm for 5 seconds. Collagen adsorption to the inner circular region of exposed glass was achieved by addition of 200 μ l of collagen I: water in 1:1 ratio, pH 5.0, and incubation at 37°C for 45 min. Wafers were then sterilized overnight in 70% ethanol in water, rinsed in water, exposed to 0.05% bovine serum albumin and rinsed with serum-free hepatocyte media (as previously described). Hepatocytes were seeded in serum-free media as previously described and allowed to spread overnight.

The following day, PET caps were applied to PDMS annuli under sterile conditions, growth-arrested (mitomycin C treatment described above) fibroblasts were seeded and allowed to attach for 1 hour, rinsed twice with 'fibroblast media', followed by removal of annuli and cap. The separated co-culture was rinsed once more

- 47 -

with fibroblast media and fibroblasts were allowed to spread for 6 hours prior to replacement of fibroblast media with 'hepatocyte media with serum'. Control co-culture was performed by methods described previously on 0.68 cm hepatocyte island patterns (as described above). Briefly, glass was modified by immobilization of collagen I, hepatocytes were seeded followed by fibroblasts. No cap or polymer annulus was applied in this condition.

Finally, absence of overlying fibroblasts on hepatocyte island was confirmed using fluorescent labels CMFDA (chloromethylfluorescein diacetate, C-2925, Molecular Probes) and CMFTR (chloromethylbenzoylaminotetramethyl rhodamine, C-2927). Cells were loaded by incubation in 25 μ M dye in media for 45 minutes, rinsed, and incubated for 30 minutes prior to a final rinse. Fibroblasts were then trypsinized as previously described and utilized in the above protocol. Separated co-cultures were rinsed and imaged 7 hours after initial fibroblast seeding.

Agitation

In order to examine the influence of fluid convection on heterogeneity in hepatocyte phenotype, co-cultures were conducted in static and 'shaken' conditions. One representative pattern was utilized for this study. Micropatterned co-cultures were generated utilizing 490 μ m hepatocyte islands with 1230 μ m center-to-center spacing as described previously. 750,000 NIH 3T3-J2 fibroblasts were added 24 hours after initial hepatocyte seeding. Replicate cultures were then cultured under two different conditions: (1) under static culture conditions as previously described and (2) under 'shaken' conditions by culturing on a rocking platform at approximately 1 Hz within a separate incubator. Media (2

- 48 -

mL) was replaced daily. Cultures were fixed and stained for intracellular albumin at indicated times.

RESULTS

Effect of Homotypic Hepatocyte Interactions on Spatial

5 Pattern of Immunostaining

We extensively probed the previously described observation that co-culture with fibroblasts induced spatial heterogeneity in intracellular albumin staining. Specifically, we examined the potential contribution of
10 homotypic hepatocyte interaction to spatial heterogeneity by studying micropatterns with different levels of homotypic interaction both in the presence and absence of fibroblasts. Patterns of intracellular albumin for five different micropatterned hepatocyte configurations were
15 compared after 48 and 144 hours of culture. Uniform distribution of intracellular albumin was detected at 48 hours in all patterns, which diminished over the time in micropatterned hepatocytes alone. Micropatterned co-cultures (i.e. addition of fibroblasts at 24 hours of
20 culture) displayed a uniform distribution of intracellular albumin similar to that observed in micropatterned hepatocyte cultures. After 6 days of co-culture, however, hepatocytes display differential levels of staining. Hepatocytes far from the heterotypic
25 interface exhibit a similar behavior to hepatocytes cultured in the absence of fibroblasts, low levels of staining. In contrast, hepatocytes proximal to the heterotypic interface exhibit relatively high levels of intracellular albumin. Thus, homotypic hepatocyte
30 interactions do not seem to be the sole contributor to the observed spatial heterogeneity in hepatocyte phenotype.

Use of Conditioned Media

In order to examine the possible induction of
35 hepatic differentiation by secreted fibroblast products,

- 49 -

experiments were conducted with hepatocytes treated with 'conditioned media'. Urea synthesis was measured as a marker of liver-specific function in a variety of such culture conditions. Media was 'conditioned' by 24 hours incubation with (1) tissue culture plastic as a control (hepatocytes + media), (2) fibroblasts alone (hepatocytes + fibroblast conditioned media), or (3) co-culture of fibroblasts and hepatocytes (hepatocytes of co-culture conditioned media). These data were compared to co-cultured fibroblasts and hepatocytes which served as a positive control for the expected level of liver-specific function (co-culture + media).

These data indicate an expected decline in liver-specific function in pure hepatocyte over the first week of culture to less than 50 $\mu\text{g/day}$. A similar decline in liver-specific function was observed in cultures treated with fibroblast conditioned media indicating insufficient concentration of humoral factors for induction of hepatic differentiation. In contrast, co-cultures of hepatocyte and fibroblasts displayed up-regulation of urea synthesis from ~60 $\mu\text{g/day}$ to ~175 $\mu\text{g/day}$ over 10 days of culture followed by stable production of urea. Some cultures were treated with co-culture conditioned media to probe for humoral factors present only when both cell types were allowed to communicate. These did not display any further induction of liver-specific function over that observed in co-culture controls indicating insufficient concentration of humoral factors for induction of hepatic differentiation (note: detection of urea in this media was due to production of urea by the co-culture utilized for conditioning media - any induction of urea synthesis in the target hepatocyte population would therefore have generated a further increase in urea production over control co-cultures).

- 50 -

Physical Separation of Cell Populations

Hepatocyte and fibroblast populations were co-cultured in the same dish yet separated by an annulus of bare glass to probe the role of labile, freely
5 secreted factors in induction of hepatic functions. Phase contrast micrographs were produced of two different initial annuli dimensions translating to two different achievable separation widths. Growth-arrested fibroblasts migrated towards the central hepatocyte
10 region at a rate of approximately 500 microns per day. After 3 days, the 1500 μm initial separation was observed to have diminished completely and cell contact occurred at the periphery of the hepatocyte island. Subsequently, cells were allowed to interact for 8 days ('contact'
15 condition). In contrast, initial cell separation of 6000 microns narrowed to 500 microns over the same time frame ('non-contact'). This experimental design allowed the examination of the role of cell proximity/cell contact in induction of hepatic functions as well as the
20 elimination of overlying fibroblasts as confirmed by fluorescent dye labeling.

Hepatocytes in the 'contact' condition exhibited an intense staining pattern in the periphery of the hepatocyte island similar to the peripheral ring of
25 staining observed in the control co-culture. In contrast, hepatocytes in the 'non-contact' condition lacked significant staining for intracellular albumin. These results indicated the importance of cell proximity (< 500 μm) for differentiation of hepatocytes.
30 Furthermore, spatial heterogeneity in hepatocyte phenotype persisted despite absence of fibroblast adhesion to surface of hepatocytes, indicating that regional differences in hepatocyte staining is not due to overlying fibroblasts.

- 51 -

Agitation of Co-Cultures

Another method of examining the potential role of secreted products by fibroblasts was the addition of fluid convection to co-cultures. Under these 'shaken' conditions, humoral factors which theoretically require a high local concentration for their bioactivity would be diluted in the bulk fluid phase and the resulting pattern of hepatocyte differentiation would differ from static conditions. In addition, agitation of culture media would allow mixing of nutrients (oxygen, glucose) and thereby alleviate potential transport limitations to the center of large hepatocyte islands.

We also measured the effect of agitation of one representative micropatterned co-culture, 490 μm , as compared to static conditions. Phase contrast micrographs demonstrate that agitation did not cause any overt fibroblast damage due to mechanical shear. In addition, low magnification, bright field images of cultures stained for intracellular albumin demonstrated no significant differences in patterns of spatial heterogeneity. The 'penetration' length of the signal for hepatocyte differentiation from the heterotypic interface did not vary significantly when compared to static cultures. These data suggested (1) spatial heterogeneity of hepatocyte phenotype in static cultures was not caused by significant nutrient limitation due to diffusional transport and (2) dilution of secreted factors by mixing did not modulate the observed pattern of spatial heterogeneity.

30 DISCUSSION

We previously reported the local induction of hepatic function in primary hepatocytes by co-cultivation with murine 3T3-J2 fibroblasts in a micropatterned configuration. The experiments summarized above probe the mechanisms by which these cells interact in detail

- 52 -

using both conventional and microfabrication techniques. These experiments focused on classification of the signal(s) broadly defined as cell-associated or freely secreted. In addition, we examined potential

5 contributors to the finite 'penetration' length of this signal leading to spatial heterogeneity in the hepatocyte phenotype. The results of these experiments are summarized below.

A Cell-Associated Signal is Implicated in Induction of

10 Hepatic Function

Experiments to classify the differentiation signal as free versus bound provided evidence that the signal(s) is cell-associated. Taken together, the results of these experiments (use of conditioned media, separation of cell
15 populations within a co-culture, and agitation of micropatterned co-cultures) point towards cell-associated molecules. Neither fibroblast conditioned media nor co-culture conditioned media were able to induce hepatocellular functions in target hepatocytes,
20 indicating the absence of a freely soluble signaling molecule. These data are supported by the general lack of induction of hepatic function using mesenchyme conditioned media or transwells by other investigators (Shimaoka et al, 1987; Morin et al, 1988; Kuri-Harcuch
25 and Mendoza-Figueroa, 1989; Donato et al, 1990).

We deduced the implausibility of a freely soluble, highly labile signal from co-cultures performed with separated cell populations. Our data indicated that cell contact (or very close proximity, $< 5 \mu\text{m}$) correlated with
30 induction of liver-specific function in hepatocytes, whereas lack of contact ($> 500 \mu\text{m}$) did not induce an observable signal as measured by immunostaining of intracellular albumin. With the exception of some unique biochemicals such as nitric oxide, other highly labile

- 53 -

signals would be expected to signal hepatocytes across 500 μm in this separated culture configuration.

We also examined morphology of hepatocytes separated from underlying fibroblasts by a 1 mm thick collagen I hydrogel and observed fibroblastic, de-differentiated morphology after a few days of culture, further indicating the lack of a freely soluble, highly labile signal (data not shown).

The potential role of freely soluble factors whose bioactivity depends on a high local concentration was also found to be minimal by the combined results of conditioned media and agitation experiments. Any soluble factor which did not induce a signal in conditioned media due to its dilution in the larger media volume, would also be diluted in agitation experiments due to fluid convection in the media. Therefore, if fluid mixing causes reduction of the concentration of some putative soluble signaling factor below its bioactive concentration, one would not expect local induction of hepatocyte function in agitation experiments. In fact, we found similar patterns of local induction of intracellular albumin in hepatocytes in micropatterned co-cultures as compared to static controls, indicating dilution of soluble factors was not a critical limitation in induction of hepatocellular function.

Analysis of the Finite Penetration Length of Differentiation Signal

With respect to the potential contributors to the spatial heterogeneity observed in the hepatocyte phenotype in co-cultures, the experiments described herein identify three potential contributors whose effects can be discounted: (i) inadequate delivery of oxygen or other nutrients to the center of hepatocyte islands, (ii) a primary homotypic effect wherein lack of hepatocyte neighbors in island periphery induced

- 54 -

up-regulation of functions, and (iii) heterogeneous signaling from fibroblasts attached to the top surface of hepatocytes.

We determined that the role of primary hepatocyte
5 homotypic interactions in induction of spatial
heterogeneity of hepatocyte phenotype was not
significant. Specifically, we found that when
hepatocytes were cultured alone, no spatial variation in
intracellular albumin was observed as a result of
10 variations in homotypic interaction. Hepatocytes in
small islands exhibited intense, uniform staining similar
to staining patterns of hepatocytes both in the periphery
and center of larger islands, followed by a spatially
uniform decline in liver-specific function at day 6. In
15 contrast, micropatterned co-cultures exhibited marked
variations in hepatocyte phenotype where hepatocytes
adjacent to the heterotypic interface expressed greater
levels of albumin, indicating that spatial heterogeneity
is not an artifact of homotypic interactions. Other,
20 preliminary results suggest that homotypic soluble
products may affect the receptivity of hepatocytes to a
differentiation signal (data not shown).

The adequacy of diffusive transport of oxygen and
other transport was determined by comparison of static
25 and agitated micropatterned co-cultures. In both cases,
a similar pattern of induction was observed at day 4,
indicating convective mixing of media did not modify
hepatocyte behavior. Finally, the contribution of
overlying fibroblasts in the observed spatial
30 heterogeneity was also determined to be minimal.
Fibroblasts were noted to adhere to the top surface of
spread hepatocytes at larger dimensions of hepatocyte
islands with dual label vital dyes and fluorescent
microscopy (data not shown); however, experiments
35 performed to separate cell populations effectively

- 55 -

prevented fibroblast attachment to the surface of hepatocytes under these conditions. Therefore, we assessed the presence of spatial heterogeneity resulting from highly characterized initial conditions. After 8
5 days of contact between cell types, intracellular albumin immunostaining indicated the presence of peripheral staining and persistence of the heterogeneous hepatocyte response. We concluded that spatial heterogeneity could not be attributed to variations in signals arising from
10 overlying fibroblasts.

In contrast, other potential contributors to spatial heterogeneity were considered and may play a significant role. These include gap junctional communication, physical penetration of fibroblast
15 processes, and diffusion of factors through 'tissue phase' as opposed to overlying "liquid phase." Homotypic gap junctions could serve to propagate the differentiation signal from the heterotypic interface inward. Indirect immunofluorescent staining (using a
20 Connexin 32 antibody acquired from Dr. David Paul, Harvard Medical School) showed presence of gap junctions in co-cultures at day 6 (data not shown). In addition, microinjection with Lucifer Yellow suggested functionality of these junctions at day 7 (data not
25 shown). These cell junctions which allow passage of molecules of less than 1200 Daltons, may play an integral role in propagation of the signal from the heterotypic interface to subsequent hepatocytes (Lowenstein et al, 1979).

30 Morphogenesis

The role of tissue reorganization on spatial heterogeneity in hepatocyte phenotype is addressed here. Notably, reorganization of cultures (both hepatocytes alone and co-cultures) was observed in smaller pattern
35 dimensions and was significantly diminished in large

- 56 -

hepatocyte islands (greater than 490 μm). In these studies, pattern configuration at later time points was perturbed by morphogenesis in the tissue - i.e. observed patterns of staining were determined not only by initial
5 pattern configuration but also by the long-term conformation adopted by the culture. For example, 100 μm islands did not display spatial heterogeneity in albumin staining, presumably because they reorganized to a pattern where all hepatocytes were proximal to the
10 heterotypic interface. In contrast, 36 μm islands reorganized to larger dimension 'cord-like' hepatic structures where some hepatocytes were a greater distance from the heterotypic interface, resulting in spatial heterogeneity of hepatocyte phenotype. Despite the
15 existence of some reorganization in these tissues, the fundamental pattern of spatial heterogeneity remained constant; hepatic structures larger than 100 μm exhibited spatial heterogeneity in hepatocyte phenotype wherein hepatocytes far from the heterotypic interface exhibited
20 low levels of intracellular albumin. Therefore, the conclusions reached above remain well founded.

SUMMARY

In these experiments, conventional culture techniques were combined with microfabricated co-cultures
25 to show that the primary signal for differentiation of hepatocytes in hepatocyte/fibroblast co-cultures is tightly fibroblast-associated. Taken together, the results of conditioned media, separated co-culture, and agitation experiments indicated a 'cell-associated'
30 signal promotes modulation (e.g., up-regulation) of liver-specific functions. The observed finite penetration of the differentiation signal may be due to gap junctional communication, 'tissue phase' diffusion of signaling molecules, and/or physical penetration of
35 fibroblast processes.

- 57 -

With respect to the implications for the design of a co-culture-based bioreactor, evidence that the signal is fibroblast-associated suggests that fibroblasts and hepatocytes must have direct contact (i.e., occupy the same compartment in a bioreactor) to produce adequate levels of liver-specific function. These co-cultures will allow the examination of candidate biochemical signals as well as spatial configurations which minimize the fraction of hepatocytes far from the heterotypic interface, creating the potential for further improvements in bulk tissue function.

REFERENCES

The publications cited herein, and other publications generally related to this invention, are referenced below.

B. Arkles, J. R. Steinmetz, J. Zazyncny, and P. Mehta, "Factors contributing to the stability of alkoxysilanes in aqueous solution," in Silicon Compounds, Register and Review, R. Anderson, G.L. Larson and C. Smith (eds.), Hüls America, NJ, 65-73, (1991) .

K. Asunoma, J.C. Gilbert, J.E. Stein, T. Takeda, and J.P. Vacanti, "Quantitation of transplanted hepatic mass necessary to cure the gunn rat model of hyperbilirubinemia," Journal of Pediatric Surgery 27:298-301, (1992).

G. Baffet, P. Loyer, D. Glaize, A. Corly, P. Etienne, and C. Guguen-Guillouzo, "Distinct effects of cell-cell communication and corticosteroids on the synthesis and distribution of cytokeratins in cultured rat hepatocytes," Journal of Cell Science 99:609-615, (1991).

- 58 -

- H. Baier and F. Bonhoeffer, "Axon guidance by gradients of a target-derived component," *Science* 25:472-475, (1992).
- A. Baird and P. Bohlen, "Fibroblast growth factors," in
5 Peptide Growth Factors and Their Receptors I, Sporn and Roberts (eds.), Springer-Verlag, NY, (1990).
- L.R. Banner and P.H. Patterson, Major changes in the expression of the mRNAs for cholinergic differentiation factor/leukemia inhibitory factor and its receptor after
10 injury to adult peripheral nerves and ganglia. *Proceedings of National Academy of Sciences* 91:7109, (1994).
- H. Baumann, G.T. Jahreis, D.N. Sauder, and A. Koj, "Human keratinocytes and monocytes release factors which
15 regulate the synthesis of major acute phase plasma proteins in hepatic cells from man, rat and mouse," *Journal of Biological Chemistry* 259:7331, (1984).
- H. Baumann and G.G. Wong, "Hepatocyte-stimulating factor III shares structural and functional identity with
20 leukemia-inhibitory factor," *Journal of Immunology* 143:1163, (1989).
- R. Bellamkonda and P. Aebischer, "Review: Tissue engineering in the nervous system," *Biotechnology and Bioengineering* 43:543-554, (1994).
- 25 K. Bergsteinsdottir, A. Kingston, R. Mirsky, K.R. Jessen, "Rat schwann cells produce interleukin-1" *Journal of Neuroimmunology* 34:15, (1991).

- 59 -

F. Berthiaume, P.V. Moghe, M. Toner, and M.L. Yarmush,
"Effect of extracellular matrix topology on cell
structure, function, and physiological responsiveness:
hepatocytes cultured in a sandwich configuration," FASEB
5 J 10:1471-1484, (1996).

S.N. Bhatia "Development of a potential bioartificial
liver: selective adhesion of hepatocytes" Master of
Science Thesis, Massachusetts Institute of Technology,
(1993).

10 S.N. Bhatia, M. Toner, R.G. Tompkins, and M.L. Yarmush,
"Selective adhesion of hepatocytes on patterned
surfaces," Annals of the New York Academy of Sciences,
745:187-209
(1994).

15 S.N. Bhatia, M.L. Yarmush, and M. Toner, "Controlling
homotypic vs heterotypic interactions by micropatterning:
co-cultures of hepatocytes and 3T3 fibroblasts," Journal
of Biomedical Materials Research, 34(2):189-199, (1997).

D.M. Bissel, D.M. Arenson, J.J. Maher, and F.J. Roll,
20 "Support of cultured hepatocytes by a laminin-rich gel:
Evidence for a functionally significant subendothelial
matrix in normal rat liver," Journal of Clinical
Investigation 79:801-812, (1987).

H. Brenner and L.J. Gaydos, "The constrained brownian
25 movement of spherical particles in cylindrical pores of
comparable radius: models of the diffusive and convective
transport of solute molecules in membranes and porous
media," Journal of Colloid Interface Science 58:312-356,
(1977).

- 60 -

S. Britland, E. Perez-Arnaud, P. Clark, B. McGinn, P. Connolly, and G. Moores, "Micropatterning proteins and synthetic peptides on solid supports: a novel application for microelectronics fabrication technology,"
5 Biotechnology Progress 8:155-160 (1992).

J.V. Boykin and J.A. Molnar, "Burn scar and skin equivalents," in Wound Healing, I.K. Cohen et al (eds.), Saunders, Philadelphia, (1992).

J.M. Calvert, "Coplanar molecular assemblies of amino-
10 and perfluorinated alkylsilanes: Characterization and geometric definition of mammalian cell adhesion and growth," Journal of American Chemical Society 114:8435-8442 (1992).

J.M. Caron, "Induction of albumin gene transcription in
15 hepatocytes by extracellular matrix proteins," Molecular Cell Biology 10:1239-1243, (1990).

S.B. Carter, "Principles of cell motility: the direction of cell movement and cancer invasion," Nature 20E:1183-1187, (1965).

20 T.M.S. Chang, "Hemoperfusion in 1981," in Hemoperfusion, Bononini and Chang (eds.), Karger, (1981).

M. Chedid, J.S. Rubin, K.G. Csaky, and S.A. Aaronson, "Regulation of keratinocyte growth factor gene expression by interleukin 1," Journal of Biological Chemistry
25 269:10753, (1994).

P. Clark, P. Connolly, and G.R. Moores, "Cell guidance by micropatterned adhesiveness in vitro," Journal of Cell Science 103:287-292, (1992).

- 61 -

P. Clark, S. Britland, and P. Connolly, "Growth cone guidance and neuron morphology on micropatterned laminin," *Journal of Cell Science*, 105:203-212 (1993).

M.G. Clemens, M. Bauer, C. Gingalewski, E. Miescher, and
5 J. Zhang. "Editorial Review: Hepatic intercellular communication in shock and inflammation," *Shock* 2(1):1-9, (1994).

B. Clement, C. Guguen-Guillouzo, J-P Campon, D. Glaise, M. Bourel, and A. Guillouzo, "Long-term co-cultures of
10 adult human hepatocytes with rat liver epithelial cells: modulation of albumin secretion and accumulation of extracellular material," *Hepatology* 4(3):373-380, (1984).

B. Clement, C. Guguen-Guillouzo, J. Grimaud, M. Bissel, and A. Guillouzo, "Effect of hydrocortisone on deposition
15 of types I and IV collagen in primary culture of rat hepatocytes," *Cellular and Molecular Biology* 34(5):449-460, (1988).

B. Clement, C. Guguen-Guillouzo, J. Campion, D. Glaise, M. Bourel, and A. Guillouzo. " Long-term co-cultures of
20 adult human hepatocytes with rat liver epithelial cells: modulation of albumin secretion and accumulation of extracellular matrix material," *Hepatology* 4(3):373-380, (1994).

C. Coers and A.L. Woolf, (eds.), *The Innervation of
25 Muscle*, C.C. Thomas, Springfield, Illinois, (1959).

J.M. Corey, B.C. Wheeler, G.J. Brewer, "Compliance of hippocampal neurons to patterned substrate networks," *Journal of Neuroscience Research*, 30:300-307 (1991).

- 62 -

- A. Corlu, B. Kneip, C. Lhadi, G. Leray, D. Glaise, G. Baffet, D. Bourel, and C. Guguen-Guillouzo, "A plasma membrane protein is involved in cell contact-mediated regulation of tissue-specific genes in adult
5 hepatocytes," *Journal of Cell Biology* 115(2):505-515, (1991).
- A. Corlu, G.P. Ilyin, N.Gerard, B. Kneip, M. Rissel, B. Jegou, and C. Guguen-Guillouzo, "Tissue distribution of liver regulating protein," *American Journal of Pathology*
10 145(3):715-727, (1994).
- R. Cotran, V. Kumar, and S. Robbins, "Diseases of Muscle," in *Robbins Pathologic Basis of Disease*, R. Cotran, V. Kumar, and S. Robbins (eds.), WB Saunders, PA, (1989).
- 15 Darnell, H. Lodish, and D. Baltimore, "The extracellular matrix serves many functions," in *Molecular Cell Biology*, J. Darnell, H. Lodish, and D. Baltimore (eds.), 904-5, W.H. Freeman and Company, New York, (1990).
- P.F. Davies, "Biology of disease: vascular cell
20 interactions with special reference to the pathogenesis of atherosclerosis," *Laboratory Investigation*, 55:5-24 (1986).
- E.T. den Braber, J.E. de Ruijter, H.T.J. Smits, L.A. Ginsel, A.F. von Recum, and J.A. Jansen, "Effect of
25 parallel surface microgrooves and surface energy on cell growth," *Journal of Biomedical Materials Research*, 29:511-518 (1995).
- F.M. De La Vega and T. Mendoza-Figueroa, "Dimethyl sulfoxide enhances lipid synthesis and secretion by

- 63 -

long-term cultures of adult rat hepatocytes," *Biochimie*
73:621-624 (1991).

A.A. Demetriou, A. Reisner, J. Sanchez, S.M. Levenson,
and A.D. Mocioni, and J.R. Chowdury, "Transplantation of
5 microcarrier-attached hepatocytes into 90% partially
hepatectomized rats," *Hepatology* 8:1006-1009, (1988).

M. Deyme, A. Baszkin, J.E. Proust, E. Perez, and M.M.
Boissonnade, "Collagen at interfaces I. In situ collagen
adsorption at solution/air and solution/polymer
10 interfaces," *Journal Biomedical Materials Research*,
20:951-962 (1986).

J. Dich, C. Vind, and N. Grunnet, "Long-term culture of
hepatocytes: effect of hormones on enzyme activities and
metabolic capacity," *Hepatology* 8:39-45, (1988).

15 C.A. Dinarello, "Interleukin-1 and interleukin-1
antagonism," *Blood* 77:1627, (1991).

M.T. Donato, M.J. Gómez-Lechón, and J.V. Castell, "Drug
metabolizing enzymes in rat hepatocytes co-cultured with
cell lines," *In Vitro Cell and Developmental Biology*,
20 26:1057-1062, (1990).

M.T. Donato, J.V. Castell, and M.J. Gomez-Lechon,
"Cytochrome P450 activities in pure and co-cultured rat
hepatocytes: effects of model inducers, " *In Vitro Cell*
and *Developmental Biology* 30A(12):825-832, (1994).

25 N.M. Le Douarin, "An experimental analysis of liver
development," *Medical Biology.*, 53:427-455 (1975).

- 64 -

J.C.Y. Dunn, M.L. Yarmush, H.G. Koebe, and R.G. Tompkins, "Hepatocyte function and extracellular matrix geometry: Long-term culture in a sandwich configuration," FASEB J, 3:174-177 (1989).

- 5 J.C.Y. Dunn, R.G. Tompkins, and M.L. Yarmush, "Long-term in vitro function of adult hepatocytes in a collagen sandwich configuration," Biotechnology Progress, 7:237-245 (1991).

- J.C.Y. Dunn, R.G. Tompkins, and M.L. Yarmush,
10 "Hepatocytes in collagen sandwich: evidence for transcriptional and translational protein," Journal of Cell Biology 116(4):1043-1053, (1992).

- J.C.Y. Dunn, J.S. Friedberg, R.G. Tompkins, and M.L. Yarmush, "Hepatocytes from rat liver perfusions:
15 physicochemical effects on polyribosome size," ASAIO Transactions 38(4): 841-845, (1992).

- J.C.Y. Dunn, "Long-term culture of differentiated hepatocyte in a collagen sandwich configuration," Doctoral Dissertation, Massachusetts Institute of
20 Technology, (1992).

P. Ekblom, "Developmentally regulated conversion of mesenchyme to epithelium," FASEB J 3:2141-2150, (1989).

- P. Ekblom, M. Ekblom, L. Fecker, G. Klein, H. Zhang, Y. Kadoya, M-L Chu, U Mayer, and R. Timpl, " Role of
25 mesenchymal nidogen for epithelial morphogenesis in vitro," Development 120:2003-2014, (1994).

M.F. Fillinger, S.E. O'Connor, R.J. Wagner, and J.L. Cronenvett, "The effect of endothelial cell coculture on

- 65 -

smooth muscle cell proliferation," Journal of Vascular Surgery, 17:1058-1068 (1993).

M.E. Fini, K.J. Strissel, M.T. Girard, J.W. Mays, W.B. Rinehart, "Interleukin 1 alpha mediates collagenase
5 synthesis," Journal of Biological Chemistry 269:11291,
(1994).

J.M. Fraslin, B. Kneip, S. Vaulont, d. Glaise, A. Muunich, and C. Guguen-Guillouzo, "Dependence of hepatocyte-specific gene expression on cell-cell
10 interactions in primary culture," EMBO J 4(10):2487-2491,
(1985).

M. Fujita, D.C. Spray, H. Choi, J.C. Saez, L.C. Rosenberg, E.L. Hertzberg, and L.M. Reid,
"Glycosaminoglycans and proteoglycans induce gap junction
15 expression and restore transcription of tissue-specific
mRNAs in primary liver cultures," Hepatology 7:S1-S9,
(1987).

R. Gebhardt and D. Mecke, "Heterogeneous distribution of glutamine synthetase among rat liver parenchymal cell in
20 situ in primary culture," EMBO J 2:567-570, (1983).

J.H. Georger, D.A. Stenger, A.S. Rudolph, J.J. Hickman, C.S. Dulcey, and T.L. Fare, "Coplanar patterns of self-assembled monolayers for selective cell adhesion and outgrowth," Thin Solid Films 210/211:716-719, (1992).

25 S.F. Gilbert, in Developmental Biology, Sinauer Associates, Sunderland, MA, 1991.

- 66 -

I.D. Goldberg and E.M. Rosen (eds.), in
Epithelial-Mesenchymal Interactions in Cancer,
Birkhauser, Verlag, Basel, (1995).

F. Goulet, C. Normand, and O. Morin, "Cellular
5 interactions promote tissue-specific function, biomatrix
deposition and junctional communication of primary
cultured hepatocytes," Hepatology, 8(5):1010-1018 (1988).

C. Guguen-Guillouzo, B. Clement, G. Baffet, C. Beaumont,
E. Morel-Chany, D. Glaize, and A. Guillouzo, "Maintenance
10 and reversibility of active albumin secretion by adult
rat hepatocytes co-cultured with another liver epithelial
cell type," Experimental Cell Research, 143:47-54 (1983).

C. Guguen-Guillouzo, "Role of homotypic and heterotypic
cell interactions in expression of specific functions by
15 cultured hepatocytes, " in Isolated and cultured
Hepatocytes A.Guillouzo and C. Guguen-Guillouzo (eds.),
John Libbery Eurotext, INSERM:259-284, (1986).

R.W. Gundersen, "Response of sensory neurites and growth
cones to patterned substrata of laminin and fibronectin
20 in vitro, " Developmental Biology, 121:423-431 (1987).

J.A. Hammarback, S.L. Palm, L.T. Furcht, and P.C.
Letourneau, "Guidance of neurite outgrowth by pathways of
substratum-adsorbed laminin," Journal of Neuroscience
Research 13:213-220, (1985).

25 J.A. Hammarback, J.B. McCarthy, S.L. Palm, L.T. Furcht,
and P.C. Letourneau, "Growth cone guidance by
substrate-bound laminin pathways is correlated with
neuron-to-pathway adhesivity," Developmental Biology
126:29-39 (1988).

- 67 -

J.A. Hammarback, and P.C. Letourneau, "Neurite extension across regions of low cell-substratum adhesivity: implications for the guidepost hypothesis of axonal pathfinding," *Developmental Biology* 117:655-671, (1986).

- 5 R.I. Handin, S.E. Lux, and T.P. Stossel, in *Blood: Principles and Practice of Hematology*, R.I. Handin, S.E. Lux, and T.P. Stossel (eds.), JB Lippincott Co, Philadelphia, (1995).

- 10 E.J. Harfenist and R.K. Murray, "Plasma proteins, immunoglobulins, and blood coagulation," in, *Harpers Biochemistry*, R.K. Murray, D.K. Granner, P.A. Mayer, and V.W. Rodwell (eds.), Appleton and Lange, CT, (1993).

- A. Harris, "Behavior of cultured cells on substrata of variable adhesiveness," *Experimental Cell Research*
15 77:285-297, (1973).

R.G. Harrison, "The cultivation of tissues in extraneous media as a method of morphogenetic study," *Anatomic Records* 6:181-193, (1912).

- E.D. Hay, and A. Zuk, "Transformations between epithelium
20 and mesenchyme: normal, pathological, and experimentally induced," *American Journal of Kidney Diseases* 26(4):678-690, (1995).

- K.E. Healy, B. Lom, and P.E. Hockberger, "Spatial distribution of mammalian cells dictated by material
25 surface chemistry," *Biotechnology and Bioengineering* 43:792-800, (1994).

D. Herzlinger, J. Qiao, D. Cohen, N. Ramakrishna, and A.M.C. Brown, "Induction of kidney epithelial

- 68 -

morphogenesis by cells expressing wnt-1," Developmental Biology 166:815-818, (1994).

T. Hirano, "Interleukin 6" in The Cytokine Handbook, Academic Press, NY, (1994).

- 5 E. Houssaint, "Differentiation of the mouse hepatic primordium.I. an analysis of tissue interactions in hepatocyte differentiation," Cell Differentiation 9:269-279, (1990).

- 10 R.O. Hynes, "Integrins: versatility, modulation, and signaling in adhesion," Cell 69:11-25, (1992).

- G.W. Ireland, P. Dopping-Hepenstal, P. Jordan, and C. O'Neill, "Effect of patterned surfaces of adhesive islands on the shape, cytoskeleton, adhesion and behavior of swiss mouse 3T3 fibroblasts," Journal of Cell Science 15 Supplement 8:19-33, (1987).

H.C. Isom, T. Secott, I. Georgoff, C. Woodworth, and J. Mumaw, "Maintenance of differentiated rat hepatocytes in primary culture, " Proceedings of National Academy of Sciences 82:3252-3256, (1985).

- 20 H.O. Jauregui, N.R. Chowdury, and J.R. Chowdury, "Use of mammalian liver cells for artificial liver support," Cell Transplantation 5(3):353-367, (1996).

- M. Kan, J.S. Huang, P.E. Mansson, H. Yasumitsu, B.Carr, W.L. McKeehan, "Heparin-binding growth factor type 1 25 (acidic fibroblast growth factor): a potential biphasic autocrine and paracrine regulator of hepatocyte regeneration," Proceedings of National Academy of Sciences 86:7432, (1989).

- 69 -

K. Kato, S. Saunders, H. Nguyen, and M. Bernfield, "Loss of cell surface syndecan-1 causes epithelia to transform into anchorage-independent mesenchyme-like cells," Molecular Biology of the Cell 6:559-576, (1995).

- 5 D. Kleinfeld, K.H. Kahler, and P.E. Hockberger, "Controlled outgrowth of dissociated neurons on patterned substrates," Journal of Neuroscience, 8(11):4098-4120 (1988).

- 10 R.J. Klebe, "Cytoscribing: A method for micropositioning cells and the construction of two- and three-dimensional synthetic tissues," Experimental Cell Research 179:362-373, (1988).

- 15 N. Koide, K. Sakguchi, Y. Koide, K. Asano, M. Kawaguchi, H. Matsushima, T. Takenami, T. Shinji, M. Mor, and T. Tsuji. "Formation of multicellular spheroids composed of adult rat hepatocytes in dishes with positively charged surfaces and under other nonadherent environments," Experimental Cell Research 186:227-235, (1990).

- 20 W. Kuri-Harcuch and T. Mendoza-Figueroa, "Cultivation of adult rat hepatocytes on 3T3 cells: expression of various liver differentiated functions," Differentiation, 41:148-157 (1989).

- 25 R. Langenbach, L. Malick, A. Tompa, C. Kuszynski, H. Freed, E. Huberman, "Maintenance of adult rat hepatocytes on c3H/10T1/2 cells," Cancer Research, 39:3509-3514 (1979).

R. Langer and J.P. Vacanti. "Tissue Engineering," Science 260:920-924, (1993).

- 70 -

M.B. Lawrence, L.V. McIntire, and S.G. Eskin, "Effect of flow on polymorphonuclear leukocyte/endothelial cell adhesion," Blood, 70:1284-1290 (1987).

M.B. Lawrence, C.W. Smith, S.G. Eskin, and L.V. McIntire,
5 "Effect on venous shear stress on CD18-mediated neutrophil adhesion to cultured endothelium," Blood, 75: 227-237 (1990).

E. LeCluyse, K. Audus, and J.H. Hochman, "Formation of extensive canalicular networks by rat hepatocytes
10 cultured in collagen-sandwich configuration," American Physiological Society, (1994).

J.S. Lee, M. Kaibara, M. Iwaki, J. Sasabe, Y. Suzuki, and M. Kusakabe, "Selective adhesion and proliferation of cells on ion-implanted polymer domains," Biomaterials,
15 14: 12 (1993).

J. Lee, R.G. Tompkins, and M.L. Yarmush, "The importance of proline on long-term hepatocyte function in a collagen sandwich configuration: regulation of protein secretion," Biotechnology and Bioengineering 40:298-305, (1992).

20 J. Lee, J.R. Morgan, R.G. Tomkins, and M.L. Yarmush, "Proline-mediated enhancement of hepatocyte function in a collagen gel sandwich culture configuration," FASEB J, 7 (1993).

P.C. Letourneau, "Cell-to-Substratum adhesion and
25 guidance of axonal elongation," Developmental Biology 44:92-101, (1975).

O. Loreal, F. Levavasseur, C. Fromaget, D. Gros, A. Guillouzo, and B. Clement, "Cooperation of ito cells and

- 71 -

hepatocytes in the deposition of an extracellular matrix in vitro," American Journal of Pathology 143(2): 538-544, 1993.

B. Lom, K.E. Healy, and P.E. Hockberger, "A versatile
5 technique for patterning biomolecules onto glass coverslips," Journal of Neuroscience Methods, 50:385-397 (1993).

W.R. Lowenstein, "Junctional intercellular communication and the control of growth," Biochemica Biophysica et Acta
10 560:1-65, (1979).

F.H. Martin, S.V. Suggs, K.E. Langley, H.S. Lu, J. Ting, K.H. Okino, C.F. Morris, I.K. McNiece, F.W. Jacobsen, and E.A. Mendiaz, "Primary structure and functional expression of rat and human stem cell factor DNAs," Cell
15 63:203, 1990.

T. Matsuda, K. Inoue, and T. Sugawara, "Development of micropatterning technology for cultured cells," ASAIO Transactions, 36: M559-M562 (1990).

T. Matsuda, T. Sugawara, and K. Inoue, "Two-dimensional
20 cell manipulation technology," ASAIO Journal, 38:M243-M247 (1992).

R. Matsuo, M. Ukida, Y. Nishikawa, N. Omori, and T. Tsuji, "The role of Kupffer cells in complement in D-galactosamine/lipopolysaccharide-induced hepatic injury
25 of rats," Acta Med Okayama 46(5):345-354, (1992).

A. Mereau, L. Grey, C. Piquet-Pellorce, J.K. Heath, "Characterization of a binding protein for leukemia

- 72 -

inhibitory factor localized in extracellular matrix,"
Journal of Cell Biology 122:713, (1993).

M. Mesnil, J. Frasnin, C. Piccoli, H. Yamasaki, and C
Guguen-Guillouzo, "Cell contact but not junctional
5 communication (dye coupling) with biliary epithelial
cells is required for hepatocytes to maintain
differentiated functions," Experimental Cell Research
173:524-533, (1987).

G.K. Michalopoulos, H.D. Cianciulli, A.R. Novotny, A.D.
10 Kligerman, S.C. Strom, and R.L. Jirtle, "Liver
regeneration studies with rat hepatocytes in primary
culture," Cancer Research 42:4673, (1982).

G.K. Michalopoulos, and M.C. DeFrances, " Liver
Regeneration," Science 276:60-66, (1997).

15 S. Miyamoto, A. Ohashi, J. Kimura, S. Tobe, and T.
Akaike, "A novel approach for toxicity sensing using
hepatocytes on a collagen-patterned plate," Sensors and
Actuators B, 13-14:196-199 (1993).

P.V. Moghe, F. Berthiaume, R.M. Ezzell, M. Toner, R.G.
20 Tompkins, and M.L. Yarmush, "Role of extracellular matrix
composition and configuration in maintenance of
hepatocyte polarity and function," Biomaterials
17:373-385, (1996).

P.V. Moghe, R.M. Ezzell, M. Toner, R.G. Tompkins, and
25 M.L. Yarmush, "Role of beta1 integrin distribution in
morphology and function of collagen-sandwiched
hepatocytes," Tissue Engineering 3:1-16, (1997).

- 73 -

- O. Morin, F. Goulet, and C. Normand, (eds.), in Liver sinusoidal endothelial cells: isolation, purification, characterization and interaction with hepatocytes, Springer International (1988).
- 5 O. Morin and C. Normand, "Long-term maintenance of hepatocyte functional activity in co-culture: Requirements for sinusoidal endothelial cells and dexamethasone," Journal of Cellular Physiology, 129:103-110 (1986).
- 10 T. Nakamura, Y. Tomitaka, and A. Ichihara, "Density dependent growth control of adult rat hepatocytes in co-culture," Journal of Biochemistry 94:1029-1035, (1983).
- 15 B.A. Naughton, "The importance of stromal cells," in Biomedical Engineering Handbook, JD Bronzino (ed.), CRC Press, (1995).
- B.A. Naughton, J.S. Roman, B. Sibanda, J.P. Weintraub, and V. Kamali. "Stereotypic culture systems for liver and bone marrow: evidence for the development of functional
20 tissue in vitro and following implantation in vivo," Biotechnology and Bioengineering 43:810-825, (1994).
- J.M. Nitsche, "Pore diffusion of nonspherical brownian particles," Ind. Eng. Chem. Res. 34: 3606-3620, (1995).
- 25 R. Nusse and H.E. Varmus, "Wnt genes," Cell 69:1073-1087, (1992).
- S.L. Nyberg, M.V. Peshwa, W.D. Payne, W-S. Hu, and F.B. Cerra, "Evolution of the bioartificial liver: the need

- 74 -

for randomized clinical trials," American Journal of Surgery 166: 512-521, (1993).

C. Oakley and D.M. Brunette, "Topographic compensation: Guidance and directed locomotion of fibroblasts on
5 grooved micromachined substrata in the absence of microtubules," Cell Motility and the Cytoskeleton, 31:45-58 (1995).

M.J. Olson, M.A. Mancini, M. A. Venkatachalam, and A.K. Roy, "Hepatocyte cytodifferentiation and cell-to-cell
10 communication" in Cell Intercommunication, W.C. De Mello (ed.), CRC Press, Inc., Boca Raton, Florida (1990).

C. O'Neill, P. Jordan, P. Riddle, and G. Ireland, "Narrow linear strips of adhesive substratum are powerful inducers of both growth and total focal contact area,"
15 Journal of Cell Science 95:577-586, (1990).

V.M. Paralkar, S. Vukicevic, A.H. Reddi, "Transforming growth factor beta type 1 binds to collagen IV of basement membrane matrix: implications for development," Developmental Biology 143:303-308, (1991).

20 E.R. Peterson, and S.M. Crain. "Regeneration and innervation in cultures of adult mammalian skeletal muscle coupled with fetal rodent spinal cord," Experimental Neurology 36:136-159 (1972).

D. Plachov, K. Chowdury, C. Walther, D Simon, J-L Guenet,
25 and P Gruss, "Pax-8 a murine paired a box gene expressed in the developing excretory system and thyroid gland," Development 110:643-651, (1990).

- 75 -

H. Popper, C.S. Davidson, C.M. Leevy, and F. Schaffner.
"The social impact of liver disease." New England Journal
of Medicine 281:1455, (1969).

G. Ramadori, J.D. Sipe, C.A. Dinarello, S.B. Mizel, and
5 H.R. Colten, "Pretranslational modulation of acute phase
hepatic protein synthesis by murine recombinant
interleukin 1 (IL-1) and purified human IL-1," Journal of
Experimental Medicine 162:930, (1985).

J.P. Ranieri, R. Bellamkonda, J. Jacob, T.G. Vargo, J.A.
10 Gardella, and P. Aebischer, "Selective neuronal cell
attachment to a covalently patterned monoamine on
fluorinated ethylene propylene films," Journal of
Biomedical Materials Research 27:917-925 (1993).

J.P. Ranieri, R. Bellamkonda, E. J. Bekos, J.A. Gardella,
15 H.J. Mathieu, L. Ruiz, and P. Aebischer, "Spatial
control of neuronal cell attachment and differentiation
on covalently patterned laminin oligopeptide substrates,
" International Journal of Developmental Neuroscience
12(8):725-735, (1994).

20 P.D. Rathgen et al, Cell 62:1105, (1990).

A.R. Rincon-Sanchez, A. Hernandez, M. de Lourdes Lopez,
T. Mendoza-Figueroa, "Synthesis and secretion of lipids
by long-term cultures of female rat hepatocytes,"
131-138, (1993).

25 V.W. Rodwell, "Catabolism of proteins and amino acid
nitrogen," in Harpers Biochemistry, R.K. Murray, D.K.
Granner, P.A. Mayer, and V.W. Rodwell (eds.), Appleton
and Lange, CT, (1993).

- 76 -

S.Rohr, D.M. Schölly, and A.G. Kléber, "Patterned growth of neonatal rat heart cells in culture, Morphological and electrophysiological characterization," Circulation Research, 68:114-130 (1991).

- 5 M. Rojkind, P.M. Novikoff, P. Greenwel, J. Rubin, L. Rojas-Valencia, A. Campos de Carvalho, R. Stocker, D. Spray, E.L. Hertzberg, and A.W. Wolkof.
"Characterization and functional studies on rat liver fat-storing cell line and freshly isolated hepatocyte
10 coculture system," American Journal of Pathology 146(6): 1995.

- E.M. Rosen, A. Joseph, L. Jin, S. Rockwell, J.A. Elias, J. Knesel, J. Wines, J. McClellan, M.J. Kluger, I.D. Goldberg, and R. Zitnik, "Regulation of scatter factor
15 production via a soluble inducing factor," Journal of Cell Biology 127:225-234, (1994).

A. Rotem, M. Toner, R.G. Tompkins, and M. L. Yarmush, "Oxygen uptake rates in cultured rat hepatocytes," Biotechnology and Bioengineering 40:1286-1291, (1992).

- 20 A. Rotem, M. Toner, S. Bhatia, B.D. Foy, R.G. Tompkins, and M.L. Yarmush, "Oxygen is a factor determining in vitro tissue assembly: effects on attachment and spreading of hepatocytes," Biotechnology and Bioengineering 43:654-660, (1994).

- 25 U.W. Rothenpieler and G. R. Dressler, "Pax-2 is required for mesenchyme to epithelium conversion during kidney development," Development 119:711-720, (1993).

J. Rozga, F. Williams, M.S. Ro, D. Neuzil, T.D. Giorgio, G. Backfisch, A.D. Moscioni, R. Hakim, and A.A.

- 77 -

- Demetriou, "Development of a bioartificial liver: properties and function of a hollow-fiber module inoculated with liver cells, " Hepatology 17:258-265, (1993).
- 5 J. Rozga, E. Morsiani, E. LePage, A.D. Moscioni, T. Giorgio, and A.A. Demetriou, "Isolated hepatocytes in a bioartificial liver: a single group view and experience," Biotechnology and Bioengineering 43:645-653, (1994).
- 10 J. Rozga, L. Podesta, E. LePage, E. Morsiani, A. Moscioni, A. Hoffman, L. Sher, F. Villamil, G. Woolf, M. McGrath, L. Kong, H. Rosen, T. Lanman, J. Vierling, L. Makowka, and A.A. Demetriou, "A bioartificial liver to treat severe acute liver failure, " Annals of Surgery 219(5):538-546 (1994).
- 15 J.S. Rubin, H. Osada, P.W. Finch, W.G. Taylor, S. Rudikoff, S.A. Aaronson, "Purification and characterization of a newly identified growth factor specific for epithelial cells," Proceedings of National Academy of Sciences 86:802, (1989).
- 20 T.W. Sadler, Langman's Medical Embryology, Williams & Wilkins, Baltimore, (1990).
- W. Schrode, D. Mecke, and R. Gebhardt, " Induction of glutamine synthetase in periportal hepacytes by cocultivation with a liver epithelial cell line,"
- 25 European Journal of Cell Biology 53:35-41, (1991).
- P.O. Seglen, "Preparation of isolated rat liver cells," Methods in Biology, 13:29-83, (1976).

- 78 -

- S. Shimaoka, T. Nakamura, A. Ichihara, "Stimulation of growth of primary cultured adult rat hepatocytes without growth factors by coculture with nonparenchymal liver cells," *Experimental Cell Research*, 172:228-242 (1987).
- 5 D.B.A Silk, and R. Williams. "Experiences in the treatment of fulminant hepatic failure by conservative therapy, charcoal hemoperfusion and polyacrylonitrile hemodialysis," *International Journal of Artificial Organs* 1:29 (1978).
- 10 R. Singhvi, G. Stephanopoulos, D.I.C. Wang, "Review: Effects of substratum morphology on cell physiology," *Biotechnology and Bioengineering*, 43:764-771 (1994a).
- R. Singhvi, A. Kumar, G.P. Lopez, G.N. Stephanopoulos, D.I.C. Wang, G.M. Whitesides, D.E. Ingber, "Engineering
15 cell shape and function," *Science*, 264:696-698 (1994b).
- J.M.W. Slack, "Theoretical Embryology," in *From Egg to Embryo, Developmental and Cell Biology Series*, Cambridge University Press, (1991).
- M.B. Sporn and A.B. Roberts, "The transforming growth
20 factor-betas" in *Peptide Growth Factors and Their Receptors I*, Sporn and Roberts, (eds.), Springer-Verlag, NY, (1990).
- A. Soekarno, B. Lom, and P.E. Hockberger, "Pathfinding by neuroblastoma cells in culture is directed by
25 preferential adhesion to positively charged surfaces," in *NeuroImage* (1993).
- E. Sonnenberg, D. Meyer, K.M. Weidner, and C. Birchmeier, "Scatter factor/hepatocyte growth factor and its

- 79 -

- receptor, the c-met tyrosine kinase, can mediate a signal exchange between mesenchyme and epithelia during mouse development," *Journal of Cell Biology* 123:223-235, (1993).
- 5 M.B. Sporn and A.B. Roberts, "The transforming growth factor-betas" in *Peptide Growth Factors and Their Receptors I*, Sporn and Roberts (eds.), Springer-Verlag, NY, (1990).
- K. Stark, S. Vainio, G. Vassilera, and A. McMahon,
10 "Epithelial transformation of metanephric mesenchyme in the developin kidney regulated by wnt-4," *Nature* 372:679-683, (1994).
- P. Stefanovich, H.W.T. Matthew, M. Toner, R.G. Tompkins, and M.L. Yarmush, "Extracorporeal plasma perfusion of
15 cultured hepatocytes: effect of intermittent perfusion on hepatocyte function and morphology," *Journal of Surgical Research* 66:57-63, (1996).
- M. Steinberg. "Reconstruction of tissue in dissociated cells," *Science* 141:401-408, (1963).
- 20 A.J. Strain, G. McGuinness, J.S. Rubin, S.A. Aaronson, "Keratinocyte growth factor and fibroblast growth factor action on DNA synthesis in rat and human hepatocytes: modulation by heparin," *Experimental Cell Research* 210:253, (1994).
- 25 D.A. Stenger, J.H. Georger, C.S. Dulcey, J.J. Hickman, A.S. Rudolph, T.B. Nielsen, S.M. McCort, and J.M. Calvert, "Coplanar molecular assemblies of amino- and perfluorinated alkylsilanes: characterization and geometric definition of mammalian cell adhesion and

- 80 -

growth," Journal of the American Chemical Society
114:8345-8442, (1992).

M. Stoker and M. Perryman, "An epithelial scatter factor
released by embryo fibroblasts," Journal of Cell Science
5 77:209-223, (1985).

M. Stoker, E. Gherardi, M. Perryman, and J. Gray,
"Scatter factor is a fibroblast-derived modulator of
epithelial cell motility," Nature 327:238-242, (1987).

N.L. Sussman, M.G. Chong, T. Koussayer, D. He, T. Shang,
10 H.H. Whisennand, and J. Kelly, "Reversal of fulminant
hepatic failure using an extracorporeal liver assist
device," Hepatology 16:60-65, (1992).

N.L. Sussman and J.H. Kelly, "The Artificial Liver,"
Scientific American: Science and Medicine May:68-77,
15 (1995).

M. Traiser, B. Diener, D. Utesch, and F. Oesch, "The gap
junctional intercellular communication is no prerequisite
for the stabilization of xenobiotic metabolizing enzyme
activities in primary rat liver parenchymal cells in
20 vitro," In Vitro Toxicology:266-273, (1995).

Trentin JJ, "Hemopoietic microenvironment, historical
perspectives, status, and projections," in Handbook of
the hemopoietic microenvironment, Tavassoli M, (ed),
Hermana Press, NJ, (1989).

25 C. Trey, D.G. Burns, and S.J. Saunders. "Treatment of
hepatic coma by exchange blood transfusion," New England
Journal of Medicine 274:473, (1966).

- 81 -

- G.A. Truskey and J.S. Pirone, "The effect of fluid shear stress upon cell adhesion to fibronectin-treated surfaces," Journal of Biomedical Materials Research 24:1333-1353, (1990).
- 5 T.Tsukui, K. Kikuchi, A. Mabuchi, T. Sudo, T. Sakamoto, G. Asano, K. Yokomuro, "Production of interleukin-1 by primary cultured parenchymal liver cells (hepatocytes)," Experimental Cell Research 210:172, (1994).
- D. Utesch, E. Molitor, K-L Platt, and F. Oesch, "
- 10 Differential stabilization of cytochrome p-450 isoenzymes in primary cultures of adult rat liver parenchymal cells," In Vitro Cell and Developmental Biology 27A:858-863, (1991).
- R.F. Valentini, T.G. Vargo, J.A. Gardello, and P.
- 15 Aebischer, " Patterned neuronal attachment and outgrowht on surface modified, electrically charged fluoropolymer substrates," Journal of Biomaterials Science Polymer Edition 5(1/2):13-36, (1993).
- C.R. Vanderberg, and E.D. Hay, "Embryonic corneal
- 20 fibroblasts transfected with E-cadherin cDNA undergo mesenchymal-epithelial transformation," Molecular Biology of the Cell 3:5a, (1992).
- Y. Vanderberghe, L. Tee, V. Rogiers, and G. Yeoh, "
- 25 Transcriptional- and post-transcriptional-dependent regulation of glutathione S-transferase expression in rat hepatocytes as a function of culture conditions," FEBS 313(2):155-159, (1992).
- S. Vukicevik, H.K. Kleinman, F.P. Luyten, A.B. Roberts, N.S. Roche, and H. Reddi, "Identification of multiple

- 82 -

active growth factors in basement membrane matrigel suggests caution in interpretation of cellular activity related to extracellular matrix components," Experimental Cell Research 202:1-8, (1992).

- 5 M. Watabe, A. Nagafuchi, S. Tsukita, and M. Takeichi,
"Induction of polarized cell-cell association and
retardation of growth by activation of the
E-cadherin-catenin adhesion system in a dispersed
carcinoma line," Journal of Cell Biology 127:247-256,
10 (1994).

D.H. Wolfle, H. Hartmann, and K Jungermann, "Induction of phosphoenolpyruvate carboxykinase by sympathetic agents in primary culture of adult rat hepatocytes," Biochemica Biophysica Research Communications 98:1084-1090, (1981).

- 15 F. Wu, M. Peshwa, F. Cerra, W. Hu, "Entrapment of hepatocyte spheroids in a hollow fiber bioreactor as a potential bioartificial liver," Tissue Engineering 1(1):29-40, (1995).

- J. Xu and R.A.F. Clark, "Extracellular matrix alters PDGF
20 regulation of fibroblast integrins," Journal of Cell Biology 132(1&2):239-249, (1996).

- M. Yamazaki, M. Tsuchida, K. Kobayashi, T. Takezawa, and Y. Mori, " A novel method to prepare size-regulated spheroids composed of human dermal fibroblasts, "
25 Biotechnology and Bioengineering 44:38-44, (1994).

KM Zsebo, D.A. Williams, and E.N. Geissler, "Stem Cell Factor is encoded at the S1 locus of the mouse and is the ligand for the c-kit tyrosine kinase receptor," Cell 63:175, (1990).

- 83 -

S.D. Zucker and J. L. Gollan, "Physiology of the Liver,"
in Bockus Gastroenterology, W.S. Haubrich and F.
Schaffner (eds.), WB Saunders, Philadelphia, (1995).

- 84 -

What is claimed is:

1. A method for producing a micropatterned co-culture containing at least two cell types, the method comprising:

5 i) providing a protein-coated substrate, wherein a protein coating the substrate defines a micropattern on the substrate;

 ii) contacting the protein-coated substrate with cells of a first cell type suspended in a first cell
10 medium under conditions such that cells of the first cell type bind the protein of the protein-coated substrate, thereby producing a micropatterned cell-coated substrate; and

 iii) contacting the micropatterned cell-coated
15 substrate with cells of a second cell type suspended in a second cell medium under conditions such that cells of the second cell type bind the substrate, thereby producing the micropatterned co-culture, wherein one of the cell media is a non-attachment medium and one of the
20 cell media is an attachment medium.

2. The method of claim 1, wherein the cells of one of the first and second cell types are hepatocytes.

3. The method of claim 1, wherein the cells of one of the first and second cell types are selected from
25 the group consisting of Kupffer cells, Ito cells, endothelial cells, and biliary ductal cells.

4. The method of claim 1, wherein the cells of one of the first and second cell types are fibroblasts.

5. The method of claim 1, wherein the cells of
30 the first cell type are hepatocytes, and the cells of the second cell type are fibroblasts.

- 85 -

6. The method of claim 1, wherein the cells one or both of the first and second cell types are selected from the group consisting of muscle cells, bone marrow cells, stromal cells, skin cells, mesenchymal cells, and
5 parenchymal tumor cells.

7. The method of claim 6, wherein the muscle cells are smooth muscle cells.

8. The method of claim 6, wherein the skin cells are keratinocytes.

10 9. The method of claim 1, wherein the non-attachment medium is a serum-free medium.

10. The method of claim 1, wherein the attachment medium comprises serum.

11. The method of claim 1, wherein the attachment
15 medium comprises at least one attachment factor selected from the group consisting of fibronectin, selectin, RGD peptides, ICAMs, E-cadherin, integrins, and antibodies that specifically bind a cell surface protein.

12. The method of claim 11, wherein the antibody
20 specifically binds a cell surface protein selected from the group consisting of integrins, ICAMs, selectins, RGD peptides, and E-cadherins.

13. The method of claim 1, wherein the protein-coated substrate comprises collagen.

25 14. The method of claim 1, wherein the protein-coated substrate comprises fibronectin, laminin, or entactin, and combinations thereof.

- 86 -

15. The method of claim 1, wherein the first and second cell types define a micropattern in which an island of cells of either the first or second cell type is surrounded by cells of either the second or first cell type, respectively.

16. The method of claim 15, wherein the island of cells is 25-1,000 μm in diameter.

17. The method of claim 16, wherein the island of cells is 30-500 μm in diameter.

18. The method of claim 17, wherein the island of cells is 100-500 μm in diameter.

19. The method of claim 1, wherein the first and second cell types define a micropattern in which an island of cells of either the first or second cell type is surrounded by cells of either the second or first cell type, respectively, and wherein at least 30% of the cells of the island of cells are within 100 μm of an interface between the island of cells and the surrounding cells.

20. The method of claim 1, wherein the substrate is selected from the group consisting of glass, polymers, and silicon substrates.

21. A co-culture of at least two cell types produced by the method of claim 1.

22. The co-culture of claim 21, wherein the co-culture comprises hepatocytes and fibroblasts.

- 87 -

23. The co-culture of claim 21, wherein the co-culture comprises a combination of cells selected from the group consisting of:

- 5 a) hepatocytes and at least one cell type selected from the group consisting of Kupffer cells, Ito cells, endothelial cells, and biliary ductal cells;
- b) endothelial cells and smooth muscle cells;
- c) mesenchymal cells and tumorigenic parenchymal cells;
- 10 d) bone marrow cells and fibroblasts; and
- e) keratinocytes and fibroblasts.

24. A method for modulating a metabolic or synthetic function of a cell of a first cell type, the method comprising:

- 15 i) providing a protein-coated substrate, wherein a protein coating the substrate defines a micropattern on the substrate;
- ii) contacting the protein-coated substrate with cells of a first cell type suspended in a first cell
20 medium under conditions such that cells of the first cell type bind the protein of the protein-coated substrate, thereby producing a micropatterned cell-coated substrate; and
- iii) contacting the micropatterned cell-coated
25 substrate with cells of a second cell type suspended in a second cell medium under conditions such that cells of the second cell type bind the substrate, thereby producing the micropatterned co-culture, wherein:
 - a) one of the cell media is a non-attachment
30 medium and one of the cell media is an attachment medium; and
 - b) the cells of the first and second cell types define a micropattern wherein cells of the second cell type surround cells of the first cell type, and at least

- 88 -

30% of the cells of the first cell type are within 100 μm of an interface between the cells of the first cell type and the cells of the second cell type,

thereby producing a micropatterned co-culture,
5 wherein a metabolic or synthetic function of a cell of the first cell type is modulated relative to cells of the first cell type in an unpatterned co-culture that comprises cells of the first and second cell types.

25. The method of claim 24, wherein modulation is
10 detected as an increase in protein production of a cell of the first cell type.

26. The method of claim 25, wherein the cells of the first cell type are hepatocytes and modulation is detected as a change in intracellular or secreted albumin
15 of a hepatocyte.

27. The method of claim 25, wherein the cells of the first cell type are hepatocytes and modulation is detected as an change in urea synthesis in a hepatocyte.

28. The method of claim 25, wherein modulation is
20 detected as an change in DNA synthesis in a cell of the first cell type.

29. The method of claim 24, wherein the co-culture comprises a combination of cells selected from the group consisting of:
25 a) hepatocytes and at least one cell type selected from the group consisting of Kupffer cells, Ito cells, endothelial cells, and biliary ductal cells;
b) endothelial cells and smooth muscle cells;
c) mesenchymal cells and tumorigenic parenchymal
30 cells;

- 89 -

- d) bone marrow cells and fibroblasts; and
- e) keratinocytes and fibroblasts.

30. The method of claim 24, wherein the co-culture comprises hepatocytes and fibroblasts.

5 31. A co-culture of cells produced according to the method of claim 24.

32. The method of claim 24, wherein the non-attachment medium is a serum-free medium.

10 33. The method of claim 24, wherein the attachment medium comprises serum.

34. The method of claim 24, wherein the protein-coated substrate comprises a protein selected from the group consisting of collagen, fibronectin, laminin, and entactin, or combinations thereof.

15 35. The method of claim 24, wherein the micropattern defined by cells of the first and second cell types comprises an island of cells of the first cell type surrounded by cells of the second cell type.

20 36. The method of claim 35, wherein the island of cells is 25-1,000 μm in diameter.

37. The method of claim 36, wherein the island of cells is 30-500 μm in diameter.

38. The method of claim 37, wherein the island of cells is 100-500 μm in diameter.

- 90 -

39. The method of claim 24, wherein the rate at which a metabolic or synthetic function is modulated in the micropatterned co-culture is increased relative to the rate at which a metabolic or synthetic function is modulated in an unpatterned co-culture.

40. The method of claim 24, wherein a metabolic or synthetic function of cells of the first cell type is modulated at least 1.5-fold in a micropatterned co-culture, relative to a metabolic or synthetic function of cells of the first cell type in an unpatterned co-culture.

41. The method of claim 40, wherein a metabolic or synthetic function of cells of the first cell type is modulated at least 5-fold in a micropatterned co-culture, relative to a metabolic or synthetic function of cells of the first cell type in an unpatterned co-culture.

42. A method for modulating a metabolic or synthetic function of a cell of a second cell type, the method comprising:

- i) providing a protein-coated substrate, wherein a protein coating the substrate defines a micropattern on the substrate;
- ii) contacting the protein-coated substrate with cells of a first cell type suspended in a first cell medium under conditions such that cells of the first cell type bind the protein of the protein-coated substrate, thereby producing a micropatterned cell-coated substrate; and
- iii) contacting the micropatterned cell-coated substrate with cells of a second cell type suspended in a second cell medium under conditions such that cells of

- 91 -

the second cell type bind the substrate, thereby producing the micropatterned co-culture, wherein:

a) one of the cell media is a non-attachment medium and one of the cell media is an attachment medium;

5 and

b) the cells of the first and second cell types define a micropattern wherein cells of the first cell type surround cells of the second cell type, and at least 30% of the cells of the second cell type are within 100 μm of an interface between the cells of the second cell type and the cells of the first cell type,

thereby producing a micropatterned co-culture, wherein a metabolic or synthetic function of a cell of the second cell type is modulated relative to cells of the second cell type in an unpatterned co-culture that comprises cells of the second and first cell types.

43. The method of claim 42, wherein the co-culture comprises a combination of cells selected from the group consisting of:

20 a) hepatocytes and at least one cell type selected from the group consisting of Kupffer cells, Ito cells, endothelial cells, and biliary ductal cells;

b) endothelial cells and smooth muscle cells;

c) mesenchymal cells and tumorigenic parenchymal cells;

d) bone marrow cells and fibroblasts;

e) keratinocytes and fibroblasts and

f) hepatocytes and fibroblasts.

44. A co-culture produced according to the method of claim 42.

- 92 -

45. The method of claim 24, wherein modulation comprises upregulation of a metabolic or synthetic function of a cell.

46. A method for producing a micropatterned
5 co-culture containing at least two cell types, the method comprising:

- i) providing a protein coated substrate wherein a protein coated substrate defines a micropattern on the substrate;
- 10 ii) contacting the protein-coated substrate with cells of a first cell type suspended in a first cell medium under conditions such that the cells of the first cell type bind the protein of the protein-coated substrate, thereby producing a micropatterned cell-coated
15 substrate; and
- iii) contacting the micropatterned cell-coated substrate with cells of a second cell type suspended in a second cell medium under conditions such that the cells of the second cell type bind to the substrate, thereby
20 producing the micropatterned co-culture, wherein the first cell type is in non-attachment medium and second cell type has natural attachment capabilities to attach it to the substrate.

25 47. The method of claim 46, wherein the cells are fibroblasts.

48. The method of claim 46, wherein the substrate is charged.

30 49. The method of claim 1, wherein said first cell type has been genetically engineered to produce a desired product, and said second cell type produces a

- 93 -

protein, and said first cell type enables said second cell type to reproduce and grow.

1/4





FIG. 1



3/4




FIG. 3




FIG. 4

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US97/08254

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C12N 11/00, 11/02, 11/08, 5/00

US CL : 435/174, 177, 180, 395, 402, 404

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/174, 177, 180, 395, 402, 404

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, CA, BIOSIS, MEDLINE

search terms: micropattern, co-culture, substrate, protein-coated, cell-coated

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X ---- Y	BHATIA et al, Controlling Cell Interactions by Micropatterning in Co-cultures: Hepatocytes and 3T3 Fibroblasts. Journal of Biomedical Materials Research, February 1997, Vol 34, No 2, pages 189-199, entire document.	1 , 2 , 4 , 5,9,10,13, 20-22,24, 30- 34, 42-44,46, 47 ----- 3,6-8,11, 12,14-19, 23,25-29, 35-41,45, 48,49

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*&*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search 11 SEPTEMBER 1997	Date of mailing of the international search report 29 SEP 1997
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized officer DAVID M. NAFF Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US97/08254

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	IRELAND et al, Effect of Patterned Surfaces of Adhesive Islands on the Shape, Cytoskeleton, Adhesion and Behaviour of Swiss Mouse 3T3 Fibroblasts. Journal of Cell Science Supplement, 1987, Vol 8, pages 19-33, entire document.	15-19, 35-38
Y	US 4,614,517 A (RUOSLAHTI et al) 30 September 1986, entire document.	11,12
A	US 5,602,029 A (MIYAMOTO) 11 February 1997, entire document.	1-49